

# I-FABP siRNA (h): sc-41239

## BACKGROUND

Fatty acid-binding proteins, designated FABPs, are a family of homologous cytoplasmic proteins that are expressed in a highly tissue-specific manner and play an integral role in the balance between lipid and carbohydrate metabolism. FABPs mediate fatty acid (FA) and/or hydrophobic ligand uptake, transport and targeting within their respective tissues. The mechanisms underlying these actions can give rise to both passive diffusional uptake and protein-mediated transmembrane transport of FAs. FABPs are expressed in adipocytes (A-FABP), brain (B-FABP), epidermis (E-FABP, also designated psoriasis-associated FABP or PA-FABP), muscle and heart (H-FABP, also designated mammary-derived growth inhibitor or MDGI), intestine (I-FABP), liver (L-FABP), myelin (M-FABP) and testis (T-FABP). Intestinal FABP (I-FABP) is an abundant cytosolic protein abundant in small intestine epithelial cells. The human gene maps to chromosome 4q26 and has a polymorphism at codon 54, which confers an alanine-encoding allele and a threonine-encoding allele. Threonine at position 54 is associated with increased fat oxidation and Insulin resistance.

## REFERENCES

1. Baier, L.J., et al. 1995. An amino acid substitution in the human intestinal fatty acid binding protein is associated with increased fatty acid binding, increased fat oxidation, and Insulin resistance. *J. Clin. Invest.* 95: 1281-1287.
2. Veerkamp, J.H. and Maatman, R.G. 1995. Cytoplasmic fatty acid-binding proteins: their structure and genes. *Prog. Lipid Res.* 34: 17-52.
3. Hotamisligil, G.S., et al. 1996. Uncoupling of obesity from Insulin resistance through a targeted mutation in aP2, the adipocyte fatty acid binding protein. *Science* 274: 1377-1379.
4. Storch, J. and Thumser, A.E. 2000. The fatty acid transport function of fatty acid-binding proteins. *Biochim. Biophys. Acta* 1486: 28-44.
5. Glatz, J.F. and Storch, J. 2001. Unravelling the significance of cellular fatty acid-binding proteins. *Curr. Opin. Lipidol.* 12: 267-274.
6. Veerkamp, J.H. and Zimmerman, A.W. 2001. Fatty acid-binding proteins of nervous tissue. *J. Mol. Neurosci.* 16: 133-142.

## CHROMOSOMAL LOCATION

Genetic locus: FABP2 (human) mapping to 4q26.

## PRODUCT

I-FABP siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see I-FABP shRNA Plasmid (h): sc-41239-SH and I-FABP shRNA (h) Lentiviral Particles: sc-41239-V as alternate gene silencing products.

For independent verification of I-FABP (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41239A, sc-41239B and sc-41239C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

I-FABP siRNA (h) is recommended for the inhibition of I-FABP expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

I-FABP (E-9): sc-374482 is recommended as a control antibody for monitoring of I-FABP gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor I-FABP gene expression knockdown using RT-PCR Primer: I-FABP (h)-PR: sc-41239-PR (20  $\mu$ l, 538 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.