

# 17 $\beta$ -HSD siRNA (m): sc-41382

## BACKGROUND

17 $\beta$ -hydroxysteroid dehydrogenase type 1 (17 $\beta$ -HSD) catalyzes the final step in the formation of estradiol and testosterone from estrone and androstenedione, respectively. Ovarian granulosa cells and breast tissue both express 17 $\beta$ -HSD. Other tissues that express 17 $\beta$ -HSD include testis, placenta, uterus, prostate and adipose tissue. 17 $\beta$ -HSD functions as a homodimer and prefers NADP(H) over NAD(H) for oxidation and reduction. The gene encoding human 17 $\beta$ -HSD maps to chromosome 17q21.2. The importance of 17 $\beta$ -HSD to estradiol production suggests the specific inhibition of 17 $\beta$ -HSD may aid in breast cancer therapy. Breast cancer patients with an amplification of 17 $\beta$ -HSD expression statistically have a worse outcome than those without. 17 $\beta$ -HSD amplification in tamoxifen-treated patients correlates to decreased breast cancer survival.

## REFERENCES

1. Luu-The, V., et al. 1990. Structure of two in tandem human 17 $\beta$ -hydroxysteroid dehydrogenase genes. *Mol. Endocrinol.* 4: 268-275.
2. Wingqvist, R., et al. 1990. The gene for 17 $\beta$ -hydroxysteroid dehydrogenase maps to human chromosome 17, bands q12-q21, and shows an RFLP with *Scal*. *Hum. Genet.* 85: 473-476.
3. Lin, S.X., et al. 1992. Subunit identity of the dimeric 17 $\beta$ -hydroxysteroid dehydrogenase from human placenta. *J. Biol. Chem.* 267: 16182-16187.
4. Poutanen, M., et al. 1993. Differential estrogen substrate specificities for transiently expressed human placental 17 $\beta$ -hydroxysteroid dehydrogenase and an endogenous enzyme expressed in cultured COS-m6 cells. *Endocrinology* 133: 2639-2644.
5. Luu-The, V., et al. 1995. Characteristics of human types 1, 2 and 3 17 $\beta$ -hydroxysteroid dehydrogenase activities: oxidation/reduction and inhibition. *J. Steroid Biochem. Mol. Biol.* 55: 581-587.
6. Vihko, P., et al. 2001. Structure and function of 17 $\beta$ -hydroxysteroid dehydrogenase type 1 and type 2. *Mol. Cell. Endocrinol.* 171: 71-76.
7. Gunnarsson, C., et al. 2003. Amplification of HSD17B1 and ERBB2 in primary breast cancer. *Oncogene* 22: 34-40.

## CHROMOSOMAL LOCATION

Genetic locus: Hsd17b1 (mouse) mapping to 11 D.

## PRODUCT

17 $\beta$ -HSD siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see 17 $\beta$ -HSD shRNA Plasmid (m): sc-41382-SH and 17 $\beta$ -HSD shRNA (m) Lentiviral Particles: sc-41382-V as alternate gene silencing products.

For independent verification of 17 $\beta$ -HSD (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41382A, sc-41382B and sc-41382C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

17 $\beta$ -HSD siRNA (m) is recommended for the inhibition of 17 $\beta$ -HSD expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor 17 $\beta$ -HSD gene expression knockdown using RT-PCR Primer: 17 $\beta$ -HSD (m)-PR: sc-41382-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.