

ADAM7 siRNA (h): sc-41404

BACKGROUND

The ADAM (a disintegrin and metalloprotease) protein family, which includes over 30 membrane-anchored, glycosylated, Zn²⁺ dependent proteases, plays a role in cell-cell and cell-matrix interface related processes, including fertilization, muscle fusion, secretion of TNF α (tumor necrosis factor α), and modulation of the neurogenic function of Notch and Delta. The ADAM proteins possess a signal-domain, a pro-domain, a metalloprotease domain, a disintegrin domain (Integrin ligand), a cysteine-rich region, an epidermal growth factor-like domain, a transmembrane domain and a cytoplasmic tail. ADAMs are expressed in a wide range of mammalian tissues and several are abundantly expressed in the male reproductive tract. ADAM7, also designated GP-83, binds wheat germ agglutinin (WGA), and is synthesized as a protein and secreted by corpus and cauda epididymus. ADAM7 conjugates to spermatozoa during their transit in human epididymus, and may be involved in sperm maturation.

REFERENCES

1. Wolfsberg, T.G., Primakoff, P., Myles, D.G. and White, J.M. 1995. ADAM, a novel family of membrane proteins containing a disintegrin and metalloprotease domain: multipotential functions in cell-cell and cell-matrix interactions. *J. Cell Biol.* 131: 275-278.
2. Stone, A.L., Kroeger, M. and Sang, Q.X. 1999. Structure-function analysis of the ADAM family of disintegrin-like and metalloproteinase-containing proteins (review). *J. Protein Chem.* 18: 447-465.
3. Primakoff, P. and Myles, D.G. 2000. The ADAM gene family: surface proteins with adhesion and protease activity. *Trends Genet.* 16: 83-87.
4. Sun, G.H., Lin, Y.C., Cha, T.L., Yu, D.S., Chang, S.Y. and Liu, H.W. 2000. Conjugation of maturation-related wheat-germ-lectin-binding proteins to caput epididymal sperm in co-cultures with corpus epididymal epithelial cells of BALB/c mouse. *Arch. Androl.* 45: 43-52.
5. Liu, H.W., Lin, Y.C., Chao, C.F., Chang, S.Y. and Sun, G.H. 2000. GP-83 and GP-39, two glycoproteins secreted by human epididymis are conjugated to spermatozoa during maturation. *Mol. Hum. Reprod.* 6: 422-428.
6. Sun, G.H., Lin, Y.C., Guo, Y.W., Chang, S.Y. and Liu, H.W. 2000. Purification of GP-83, a glycoprotein secreted by the human epididymis and conjugated to mature spermatozoa. *Mol. Hum. Reprod.* 6: 429-434.
7. Lin, Y.C., Sun, G.H., Lee, Y.M., Guo, Y.W. and Liu, H.W. 2001. Cloning and characterization of a complementary DNA encoding a human epididymis-associated disintegrin and metalloprotease 7 protein. *Biol. Reprod.* 65: 944-950.
8. Bates, E.E., Fridman, W.H. and Mueller, C.G. 2002. The ADAMDEC1 (decysin) gene structure: evolution by duplication in a metalloprotease gene cluster on chromosome 8p12. *Immunogenetics* 54: 96-105.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: ADAM7 (human) mapping to 8p21.2.

PRODUCT

ADAM7 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ADAM7 shRNA Plasmid (h): sc-41404-SH and ADAM7 shRNA (h) Lentiviral Particles: sc-41404-V as alternate gene silencing products.

For independent verification of ADAM7 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41404A, sc-41404B and sc-41404C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ADAM7 siRNA (h) is recommended for the inhibition of ADAM7 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ADAM7 gene expression knockdown using RT-PCR Primer: ADAM7 (h)-PR: sc-41404-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.