

NPC1 siRNA (m): sc-41589

BACKGROUND

Cells obtain cholesterol via two distinct pathways, endogenous synthesis in the endoplasmic reticulum and exogenous uptake through the low-density lipoprotein (LDL) receptor pathway. NPC1 is a protein that resides in late endosomes and lysosomes and is involved in the intracellular trafficking of cholesterol. The human NPC1 gene maps to chromosome 18q11.2 and produces proteins which undergo N-glycosylation and are expressed in brain and liver. NPC1 contains a cysteine-rich domain, which is critical for proper protein function, but is highly mutated. Mutations in NPC1 result in Niemann-Pick disease type C (NPC), an autosomal recessive disease characterized by the accumulation of unesterified cholesterol in the endosomal/lysosomal system. The accumulation of cholesterol results in progressive neurodegeneration and death. More than 90% of cases of NPC are due to mutations in NPC1 and patients with NPC display multiple neurological symptoms, such as hepatosplenomegaly, ataxia, dystonia and dementia.

REFERENCES

1. Watari, H., et al. 1999. Mutations in the leucine zipper motif and sterol-sensing domain inactivate the Niemann-Pick C1 glycoprotein. *J. Biol. Chem.* 274: 21861-21866.
2. Greer, W.L., et al. 1999. Mutations in NPC1 highlight a conserved NPC1-specific cysteine-rich domain. *Am. J. Hum. Genet.* 65: 1252-1260.
3. Sym, M., et al. 2000. A model for Niemann-Pick type C disease in the nematode *Caenorhabditis elegans*. *Curr. Biol.* 10: 527-530.
4. Cruz, J.C. and Chang, T.Y. 2000. Fate of endogenously synthesized cholesterol in Niemann-Pick type C1 cells. *J. Biol. Chem.* 275: 41309-41316.

CHROMOSOMAL LOCATION

Genetic locus: *Npc1* (mouse) mapping to 18 A1.

PRODUCT

NPC1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NPC1 shRNA Plasmid (m): sc-41589-SH and NPC1 shRNA (m) Lentiviral Particles: sc-41589-V as alternate gene silencing products.

For independent verification of NPC1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41589A, sc-41589B and sc-41589C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NPC1 siRNA (m) is recommended for the inhibition of NPC1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

NPC1 (G-1): sc-271335 is recommended as a control antibody for monitoring of NPC1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NPC1 gene expression knockdown using RT-PCR Primer: NPC1 (m)-PR: sc-41589-PR (20 μ l, 575 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Xu, X., et al. 2017. Simvastatin promotes NPC1-mediated free cholesterol efflux from lysosomes through CYP7A1/LXR α signalling pathway in oxLDL-loaded macrophages. *J. Cell. Mol. Med.* 21: 364-374.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.