

G_α olf siRNA (h): sc-41754

BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (a photon, pheromone, odorant, hormone or neurotransmitter) while the effectors (e.g. adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. In mammals, G protein α , β and γ polypeptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their α subunits, since these proteins bind and hydrolyze GTP, and most obviously regulate the activity of the best studied effectors. The G_s subfamily of G α subunits includes two closely related proteins, G α_s and G α_{olf} , which, respectively, stimulate adenylyl cyclase and mediate response to olfactory stimuli.

REFERENCES

1. Jones, D.T., et al. 1990. Biochemical characterization of three stimulatory GTP-binding proteins. The large and small forms of G_s and the olfactory-specific G protein, G_{olf}. J. Biol. Chem. 265: 2671-2676.
2. Simon, M.I., et al. 1991. Diversity of G proteins in signal transduction. Science 252: 802-808.
3. Iñíguez-Lluhi, J.A., et al. 1992. G protein $\beta\gamma$ subunits synthesized in Sf9 cells. J. Biol. Chem. 267: 23409-23417.
4. Cali, J.J., et al. 1992. Selective tissue distribution of G protein γ subunits, including a new form of the γ subunits identified by cDNA cloning. J. Biol. Chem. 267: 24023-24027.
5. McLaughlin, S.K., et al. 1992. Gustducin is a taste-cell-specific G protein closely related to the transducins. Nature 357: 563-569.
6. von Weizsäcker, E., et al. 1992. Diversity among the β subunits of heterotrimeric GTP-binding proteins: characterization of a novel β subunit cDNA. Biochem. Biophys. Res. Commun. 183: 350-356.
7. Kleuss, C., et al. 1992. Different β subunits determine G protein interaction with transmembrane receptors. Nature 358: 424-426.
8. Conklin, B.R. and Bourne, H.R. 1993. Structural elements of G α subunits that interact with G $\beta\gamma$ receptors, and effectors. Cell 73: 631-641.
9. Kleuss, C., et al. 1993. Selectivity in signal transduction determined by γ subunits of heterotrimeric G proteins. Science 259: 832-834.

CHROMOSOMAL LOCATION

Genetic locus: GNAL (human) mapping to 18p11.21.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

G_α olf siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see G_α olf shRNA Plasmid (h): sc-41754-SH and G_α olf shRNA (h) Lentiviral Particles: sc-41754-V as alternate gene silencing products.

For independent verification of G_α olf (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41754A, sc-41754B and sc-41754C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

G_α olf siRNA (h) is recommended for the inhibition of G_α olf expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor G_α olf gene expression knockdown using RT-PCR Primer: G_α olf (h)-PR: sc-41754-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.