

Rab 6C siRNA (h): sc-41826

BACKGROUND

The Ras-related superfamily of guanine nucleotide binding proteins includes the R-Ras, Rap, Ral/Rec and Rho/Rab subfamilies. Increasing data suggests an important role for Rab proteins in either endocytosis or in biosynthetic protein transport. The process of transporting newly synthesized proteins from the endoplasmic reticulum to various stacks of the Golgi complex and to secretory vesicles involves the movement of carrier vesicles and requires Rab protein function. Rab proteins are also an integral part of endocytic pathways. Rab 6B, a 208 amino acid protein, and Rab 6C (also known as WTH3), a 254 amino acid protein, are members of the Rab family of guanine nucleotide binding proteins. Localized to the Golgi apparatus, Rab 6B is predominately expressed in brain tissue where it is thought to play a role in retrograde membrane traffic within the Golgi. While Rab 6B is involved in protein transport events, Rab 6C interacts with p53 and functions to negatively regulate multidrug resistance (MDR), possibly contributing to drug resistance in breast cancer cells.

REFERENCES

1. Chen, D., Guo, J. and Gahl, W.A. 1997. Rab GTPases expressed in human melanoma cells. *Biochim. Biophys. Acta* 1355: 1-6.
2. Fitzgerald, M.L. and Reed, G.L. 1999. Rab 6 is phosphorylated in thrombin-activated platelets by a protein kinase C-dependent mechanism: effects on GTP/GDP binding and cellular distribution. *Biochem. J.* 342: 353-360.
3. Shan, J., Mason, J.M., Yuan, L., Barcia, M., Porti, D., Calabro, A., Budman, D., Vinciguerra, V. and Xu, H. 2000. Rab 6C, a new member of the Rab gene family, is involved in drug resistance in MCF7/AdrR cells. *Gene* 257: 67-75.
4. Opdam, F.J., Echard, A., Croes, H.J., van den Hurk, J.A., van de Vorstenbosch, R.A., Ginsel, L.A., Goud, B. and Fransen, J.A. 2000. The small GTPase Rab 6B, a novel Rab 6 subfamily member, is cell-type specifically expressed and localised to the Golgi apparatus. *J. Cell Sci.* 113: 2725-2735.
5. Shan, J., Yuan, L., Budman, D.R. and Xu, H.P. 2002. WTH3, a new member of the Rab 6 gene family, and multidrug resistance. *Biochim. Biophys. Acta* 1589: 112-123.
6. Tian, K., Jurukovski, V., Wang, X.P., Kaplan, M.H. and Xu, H. 2005. Epigenetic regulation of WTH3 in primary and cultured drug-resistant breast cancer cells. *Cancer Res.* 65: 10024-10031.
7. Garcia-Saez, I., Tcherniuk, S. and Kozielski, F. 2006. The structure of human neuronal Rab 6B in the active and inactive form. *Acta Crystallogr. D Biol. Crystallogr.* 62: 725-733.
8. Tian, K., Wang, Y. and Xu, H. 2007. WTH3 is a direct target of the p53 protein. *Br. J. Cancer* 96: 1579-1586.
9. Wanschers, B.F., van de Vorstenbosch, R., Schlager, M.A., Splinter, D., Akhmanova, A., Hoogenraad, C.C., Wieringa, B. and Fransen, J.A. 2007. A role for the Rab 6B Bicaudal-D1 interaction in retrograde transport in neuronal cells. *Exp. Cell Res.* 313: 3408-3420.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: RAB6C (human) mapping to 2q21.1.

PRODUCT

Rab 6C siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rab 6C shRNA Plasmid (h): sc-41826-SH and Rab 6C shRNA (h) Lentiviral Particles: sc-41826-V as alternate gene silencing products.

For independent verification of Rab 6C (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41826A, sc-41826B and sc-41826C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rab 6C siRNA (h) is recommended for the inhibition of Rab 6C expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rab 6C gene expression knockdown using RT-PCR Primer: Rab 6C (h)-PR: sc-41826-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.