

Neurogenin 2 siRNA (h): sc-42077

BACKGROUND

The Neurogenin family of proteins belongs to the basic helix-loop-helix (bHLH) superfamily and consists of Neurogenin 1, Neurogenin 2 and Neurogenin 3 (also designated *ngn3*). bHLH members are transcriptional regulators that determine cell fate. During mouse neurogenesis, Neurogenin 1 and Neurogenin 2 are expressed in distinct progenitor populations in the central and peripheral nervous systems. Targeted mutation analyses showed that Neurogenin 1 is essential for the determination of neuronal precursors for proximal cranial sensory ganglia and that Neurogenin 2 is essential for the determination of precursors for epibranchial placode-derived sensory neurons.

REFERENCES

1. Ben-Arie, N., et al. 1996. Evolutionary conservation of sequence and expression of the bHLH protein Atonal suggests a conserved role in neurogenesis. *Hum. Mol. Genet.* 5: 1207-1216.
2. Tamimi, R.M., et al. 1997. *NEUROD2* and *NEUROD3* genes map to human chromosomes 17q12 and 5q23-q31 and mouse chromosomes 11 and 13, respectively. *Genomics* 40: 355-357.
3. Ma, Q., et al. 1998. Neurogenin 1 is essential for the determination of neuronal precursors for proximal cranial sensory ganglia. *Neuron* 20: 469-482.
4. Fode, C., et al. 1998. The bHLH protein *NEUROGENIN 2* is a determination factor for epibranchial placode-derived sensory neurons. *Neuron* 20: 483-494.
5. Jacquemin, P., et al. 2000. Transcription factor hepatocyte nuclear factor 6 regulates pancreatic endocrine cell differentiation and controls expression of the proendocrine gene *ngn3*. *Mol. Cell. Biol.* 20: 4445-4454.
6. Gradwohl, G., et al. 2000. Neurogenin 3 is required for the development of the four endocrine cell lineages of the pancreas. *Proc. Natl. Acad. Sci. USA* 97: 1607-1611.
7. Schwitzgebel, V.M., et al. 2000. Expression of Neurogenin 3 reveals an islet cell precursor population in the pancreas. *Development* 127: 3533-3542.
8. Jensen, J., et al. 2000. Independent development of pancreatic α - and β -cells from Neurogenin 3-expressing precursors: a role for the notch pathway in repression of premature differentiation. *Diabetes* 49: 163-176.

CHROMOSOMAL LOCATION

Genetic locus: *NEUROG2* (human) mapping to 4q25.

PRODUCT

Neurogenin 2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Neurogenin 2 shRNA Plasmid (h): sc-42077-SH and Neurogenin 2 shRNA (h) Lentiviral Particles: sc-42077-V as alternate gene silencing products.

For independent verification of Neurogenin 2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42077A, sc-42077B and sc-42077C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Neurogenin 2 siRNA (h) is recommended for the inhibition of Neurogenin 2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Neurogenin 2 (2A8): sc-293430 is recommended as a control antibody for monitoring of Neurogenin 2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Neurogenin 2 gene expression knockdown using RT-PCR Primer: Neurogenin 2 (h)-PR: sc-42077-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.