



VDAC3 siRNA (h): sc-42359

BACKGROUND

Adenine nucleotide translocator (ANT) and the voltage-dependent anion-selective channel proteins 1, 2 and 3 (VDAC1, VDAC2 and VDAC3) are components of the permeability transition pore complex (PTPC) of the mitochondrial inner or outer membranes. Formation of PTPCs, the subsequent dissipation of mitochondrial inner membrane potential and release of cytochrome c through the outer mitochondrial membrane are critical events in the early stages of apoptosis. Bax, a proapoptotic protein, has been shown to act upon ANT to induce the dissipation of mitochondrial inner membrane potential.

REFERENCES

1. Cozens, A.L., et al. 1989. DNA sequences of two expressed nuclear genes for human mitochondrial ADP/ATP translocase. *J. Mol. Biol.* 206: 261-280.
2. Li, K., et al. 1989. A human muscle adenine nucleotide translocator gene has four exons, is located on chromosome 4, and is differentially expressed. *J. Biol. Chem.* 264: 13998-14004.
3. Blachly-Dyson, E., et al. 1993. Cloning and functional expression in yeast of two human isoforms of the outer mitochondrial membrane channel, the voltage-dependent anion channel. *J. Biol. Chem.* 268: 1835-1841.
4. Zamzami, N., et al. 1996. Mitochondrial control of nuclear apoptosis. *J. Exp. Med.* 183: 1533-1544.
5. Green, D.R., et al. 1998. Mitochondria and apoptosis. *Science* 281: 1309-1312.
6. Marzo, I., et al. 1998. Bax and adenine nucleotide translocator cooperate in the mitochondrial control of apoptosis. *Science* 281: 2027-2031.
7. Kumarswamy, R. and Chandna, S. 2009. Putative partners in Bax mediated cytochrome c release: ANT, CypD, VDAC or none of them? *Mitochondrion* 9: 1-8.
8. Qin, G., et al. 2009. Oxidative damage of mitochondrial proteins contributes to fruit senescence: a redox proteomics analysis. *J. Proteome Res.* 8: 2449-2462.
9. Halestrap, A.P. 2009. What is the mitochondrial permeability transition pore? *J. Mol. Cell. Cardiol.* 46: 821-831.

CHROMOSOMAL LOCATION

Genetic locus: VDAC3 (human) mapping to 8p11.21.

PRODUCT

VDAC3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see VDAC3 shRNA Plasmid (h): sc-42359-SH and VDAC3 shRNA (h) Lentiviral Particles: sc-42359-V as alternate gene silencing products.

For independent verification of VDAC3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42359A, sc-42359B and sc-42359C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

VDAC3 siRNA (h) is recommended for the inhibition of VDAC3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor VDAC3 gene expression knockdown using RT-PCR Primer: VDAC3 (h)-PR: sc-42359-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.