

RyR-3 siRNA (m): sc-42639

BACKGROUND

Dihydropyridine receptor (DHPR) and the sarcoplasmic reticulum ryanodine receptor (RyR) are two key components of the intracellular junctions, where depolarization of the surface membrane is converted into the release of Ca^{2+} from internal stores. The RyR family consists of RyR-1, RyR-2 and RyR-3 which are characterized respectively as skeletal muscle, cardiac and brain ryanodine receptors. RyR proteins are essential for calcium-dependent excitation. Cells that do not express RyR lack excitation-contraction coupling and exhibit a several-fold reduction in Ca^{2+} current density. RyR-3 is expressed in the corpus striatum, thalamus and hippocampus as well as smooth muscle. In response to β -Amyloid, RyR-3 levels are increased, altering Ca^{2+} homeostasis in neurons. RyR-3 may play a significant role in the progression of Alzheimer's disease.

REFERENCES

1. Otsu, K., et al. 1990. Molecular cloning of cDNA encoding the Ca^{2+} release channel (ryanodine receptor) of rabbit cardiac muscle sarcoplasmic reticulum. *J. Biol. Chem.* 265: 13472-13483.
2. Hakamata, Y., et al. 1992. Primary structure and distribution of a novel ryanodine receptor/calcium release channel from rabbit brain. *FEBS Lett.* 312: 229-235.
3. Otsu, K., et al. 1993. Chromosome mapping of five human cardiac and skeletal muscle sarcoplasmic reticulum protein genes. *Genomics* 17: 507-509.
4. Sorrentino, V., et al. 1993. Localization of a novel ryanodine receptor gene (RyR3) to human chromosome 15q14-q15 by *in situ* hybridization. *Genomics* 18: 163-165.
5. Flucher, B.E., et al. 1996. Formation of junctions involved in excitation-contraction coupling in skeletal and cardiac muscle. *Proc. Natl. Acad. Sci. USA* 93: 8101-8106.
6. Franzini-Armstrong, C., et al. 1997. Ryanodine receptors of striated muscles: a complex channel capable of multiple interactions. *Physiol. Rev.* 77: 699-729.
7. Nakai, J., et al. 1997. Functional nonequality of the cardiac and skeletal ryanodine receptors. *Proc. Natl. Acad. Sci. USA* 94: 1019-1022.

CHROMOSOMAL LOCATION

Genetic locus: RyR3 (mouse) mapping to 2 E3.

PRODUCT

RyR-3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RyR-3 shRNA Plasmid (m): sc-42639-SH and RyR-3 shRNA (m) Lentiviral Particles: sc-42639-V as alternate gene silencing products.

For independent verification of RyR-3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42639A, sc-42639B and sc-42639C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RyR-3 siRNA (m) is recommended for the inhibition of RyR-3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

RyR (F-1): sc-376507 is recommended as a control antibody for monitoring of RyR-3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RyR-3 gene expression knockdown using RT-PCR Primer: RyR-3 (m)-PR: sc-42639-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.