

# L-type Ca<sup>++</sup> CP α1F siRNA (h): sc-42692

## BACKGROUND

Voltage-dependent Ca<sup>2+</sup> channels mediate Ca<sup>2+</sup> entry into excitable cells in response to membrane depolarization, and they are involved in a variety of Ca<sup>2+</sup>-dependent processes, including muscle contraction, hormone or neurotransmitter release and gene expression. Calcium channels are highly diverse, multimeric complexes composed of an α-1 subunit, an intracellular β-subunit, a disulfide linked α-2/δ subunit and a transmembrane γ-subunit. Ca<sup>2+</sup> currents are characterized on the basis of their biophysical and pharmacologic properties and include L-, N-, T-, P-, Q-, and R- types. L-type Ca<sup>2+</sup> currents initiate muscle contraction, endocrine secretion, and gene transcription, and can be regulated through second-messenger activated protein phosphorylation pathways. L-type calcium channels may form macromolecular signaling complexes with G protein-coupled receptors, thereby enhancing the selectivity of regulating specific targets.

## REFERENCES

1. Perez-Reyes, E., et al. 1995. Molecular biology of calcium channels. *Kidney Int.* 48: 1111-1124.
2. Randall, A.D. 1998. The molecular basis of voltage-gated Ca<sup>2+</sup> channel diversity: is it time for T? *J. Membr. Biol.* 161: 207-213.
3. Catterall, W.A. 2000. Structure and regulation of voltage-gated Ca<sup>2+</sup> channels. *Annu. Rev. Cell Dev. Biol.* 16: 521-555.
4. Davare, M.A., et al. 2001. A β<sub>2</sub> adrenergic receptor signaling complex assembled with the Ca<sup>2+</sup> channel Ca<sub>v</sub>1.2. *Science* 293: 98-101.
5. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 601011. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: CACNA1F (human) mapping to Xp11.23.

## PRODUCT

L-type Ca<sup>++</sup> CP α1F siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see L-type Ca<sup>++</sup> CP α1F shRNA Plasmid (h): sc-42692-SH and L-type Ca<sup>++</sup> CP α1F shRNA (h) Lentiviral Particles: sc-42692-V as alternate gene silencing products.

For independent verification of L-type Ca<sup>++</sup> CP α1F (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42692A, sc-42692B and sc-42692C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

L-type Ca<sup>++</sup> CP α1F siRNA (h) is recommended for the inhibition of L-type Ca<sup>++</sup> CP α1F expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

L-type Ca<sup>++</sup> CP α1F (1H6): sc-517005 is recommended as a control antibody for monitoring of L-type Ca<sup>++</sup> CP α1F gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor L-type Ca<sup>++</sup> CP α1F gene expression knockdown using RT-PCR Primer: L-type Ca<sup>++</sup> CP α1F (h)-PR: sc-42692-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.