

HLA-DO β siRNA (m): sc-42914

BACKGROUND

Peptide (antigen) binding to major histocompatibility complex (MHC) class II molecules destined for presentation to CD4⁺ helper T cells is determined by two key events. These include the dissociation of class II-associated invariant chain peptides (CLIP) from an antigen binding groove in MHC II-Ig dimers and by the activity of MHC molecules HLA-DM and -DO. Accumulating in endosomal/lysosomal compartments and on the surface of B cells, HLA-DM, -DO molecules regulate the dissociation of CLIP and the subsequent binding of exogenous peptides to HLA class II molecules (HLA-DR) by sustaining a conformation that favors peptide exchange. RFLP analysis of HLA-DM genes from rheumatoid arthritis (RA) patients suggests that certain polymorphisms are genetic factors for RA susceptibility.

REFERENCES

1. Kropshofer, H., Vogt, A.B., Thery, C., Armandola, E.A., Li, B.C., Moldenhauer, G., Amigorena, S. and Hammerling, G.J. 1998. A role for HLA-DO as a co-chaperone of HLA-DM in peptide loading of MHC class II molecules. *EMBO J.* 17: 2971-2981.
2. Siegmund, T., Donner, H., Braun, J., Usadel, K.H. and Badenhoop, K. 1999. HLA-DM α and HLA-DM β alleles in German patients with type 1 diabetes mellitus. *Tissue Antigens* 54: 291-294.
3. Arndt, S.O., Vogt, A.B., Markovic-Plese, S., Martin, R., Moldenhauer, G., Wolpi, A., Sun, Y., Schadendorf, D., Hammerling, G.J. and Kropshofer, H. 2000. Functional HLA-DM on the surface of B cells and immature dendritic cells. *EMBO J.* 19: 1241-1251.
4. Brunet, A., Samaan, A., Deshaies, F., Kindt, T.J. and Thibodeau, J. 2000. Functional characterization of a lysosomal sorting motif in the cytoplasmic tail of HLA-DO β . *J. Biol. Chem.* 275: 37062-37071.
5. Doebele, C.R., Busch, R., Scott, M.H., Pashine, A. and Mellins, D.E. 2000. Determination of the HLA-DM interaction site on HLA-DR molecules. *Immunity* 13: 517-527.
6. Louis-Pence, P., Kerlan-Candon, S., Morel, J., Combe, B., Clot, J., Pinet, V. and Eliaou, J.F. 2000. The down-regulation of HLA-DM gene expression in rheumatoid arthritis is not related to their promoter polymorphism. *J. Immunol.* 165: 4861-4869.
7. Toussiot, E., Sauvageot, C., Chabod, J., Ferrand, C., Tiberghien, P. and Wendling, D. 2000. The association of HLA-DM genes with rheumatoid arthritis in eastern France. *Hum. Immunol.* 61: 303-308.

CHROMOSOMAL LOCATION

Genetic locus: H2-Ob (mouse) mapping to 17 B1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

HLA-DO β siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HLA-DO β shRNA Plasmid (m): sc-42914-SH and HLA-DO β shRNA (m) Lentiviral Particles: sc-42914-V as alternate gene silencing products.

For independent verification of HLA-DO β (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42914A, sc-42914B and sc-42914C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HLA-DO β siRNA (m) is recommended for the inhibition of HLA-DO β expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HLA-DO β gene expression knockdown using RT-PCR Primer: HLA-DO β (m)-PR: sc-42914-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.