

# TCL-1B3 siRNA (m): sc-42993

## BACKGROUND

T cell leukemia/lymphoma protein 1B, TCL-1B, (also designated TML1, TCL/MTCP1-like 1, syncytiotrophoblast-specific protein and SYN-1), is a protein involved in T cell prolymphocytic leukemia (T-PLL). TCL-1B is located within the region on chromosome 14q32.1 which is frequently involved in chromosomal translocations and inversions with one of the T cell receptor loci in human T cell leukemias and lymphomas. TCL-1B is activated by chromosomal rearrangements involving the TCL1 locus. In mouse, TCL-1B is represented by five homologues, TCL-1B1–TCL-1B5. The crystal structure of TCL-1B suggests that it may play a role in the transport of small molecules such as retinoids, nucleotides and fatty acids. TCL-1B is found in both the nucleus and the cytoplasm of normal bone marrow and peripheral lymphocytes.

## REFERENCES

1. Fu, T.B., Virgilio, L., Narducci, M.G., Facchiano, A., Russo, G. and Croce, C.M. 1994. Characterization and localization of the TCL-1 oncogene product. *Cancer Res.* 54: 6297-6301.
2. Fu, Z.Q., Du Bois, G.C., Song, S.P., Kulikovskaya, I., Virgilio, L., Rothstein, J.L., Croce, C.M., Weber, I.T. and Harrison, R.W. 1998. Crystal structure of MTCP-1: implications for role of TCL-1 and MTCP-1 in T cell malignancies. *Proc. Natl. Acad. Sci. USA* 95: 3413-3418.
3. Sugimoto, J., Hatakeyama, T., Narducci, M.G., Russo, G. and Isobe, M. 1999. Identification of the TCL-1/MTCP-1-like 1 (TML1) gene from the region next to the TCL-1 locus. *Cancer Res.* 59: 2313-2317.
4. Pekarsky, Y., Hallas, C., Isobe, M., Russo, G. and Croce, C.M. 1999. Abnormalities at 14q32.1 in T cell malignancies involve two oncogenes. *Proc. Natl. Acad. Sci. USA* 96: 2949-2951.
5. Hallas, C., Pekarsky, Y., Itoyama, T., Varnum, J., Bichi, R., Rothstein, J. and Croce, C.M. 1999. Genomic analysis of human and mouse TCL1 loci reveals a complex of tightly clustered genes. *Proc. Natl. Acad. Sci. USA* 96: 14418-14423.
6. Pekarsky, Y., Koval, A., Hallas, C., Bichi, R., Tresini, M., Malstrom, S., Russo, G., Tsichlis, P. and Croce, C.M. 2000. TCL-1 enhances Akt kinase activity and mediates its nuclear translocation. *Proc. Natl. Acad. Sci. USA* 97: 3028-3033.

## CHROMOSOMAL LOCATION

Genetic locus: Tcl1b5 (mouse) mapping to 12 E.

## PRODUCT

TCL-1B3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCL-1B3 shRNA Plasmid (m): sc-42993-SH and TCL-1B3 shRNA (m) Lentiviral Particles: sc-42993-V as alternate gene silencing products.

For independent verification of TCL-1B3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42993A, sc-42993B and sc-42993C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

TCL-1B3 siRNA (m) is recommended for the inhibition of TCL-1B3 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCL-1B3 gene expression knockdown using RT-PCR Primer: TCL-1B3 (m)-PR: sc-42993-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.