

TCL-1B4 siRNA (m): sc-42995

BACKGROUND

T cell leukemia/lymphoma protein 1B, TCL-1B, (also designated TML1, TCL/MTCP1-like 1, syncytiotrophoblast-specific protein and SYN-1), is a protein involved in T cell prolymphocytic leukemia (T-PLL). TCL-1B is located within the region on chromosome 14q32.1 which is frequently involved in chromosomal translocations and inversions with one of the T cell receptor loci in human T cell leukemias and lymphomas. TCL-1B is activated by chromosomal rearrangements involving the TCL1 locus. In mouse, TCL-1B is represented by five homologues, TCL-1B1–TCL-1B5. The crystal structure of TCL-1B suggests that it may play a role in the transport of small molecules such as retinoids, nucleotides and fatty acids. TCL-1B is found in both the nucleus and the cytoplasm of normal bone marrow and peripheral lymphocytes.

REFERENCES

1. Fu, T.B., Virgilio, L., Narducci, M.G., Facchiano, A., Russo, G. and Croce, C.M. 1994. Characterization and localization of the TCL-1 oncogene product. *Cancer Res.* 54: 6297-6301.
2. Fu, Z.Q., Du Bois, G.C., Song, S.P., Kulikovskaya, I., Virgilio, L., Rothstein, J.L., Croce, C.M., Weber, I.T. and Harrison, R.W. 1998. Crystal structure of MTCP-1: implications for role of TCL-1 and MTCP-1 in T cell malignancies. *Proc. Natl. Acad. Sci. USA* 95: 3413-3418.
3. Sugimoto, J., Hatakeyama, T., Narducci, M.G., Russo, G. and Isobe, M. 1999. Identification of the TCL-1/MTCP-1-like 1 (TML1) gene from the region next to the TCL-1 locus. *Cancer Res.* 59: 2313-2317.
4. Pekarsky, Y., Hallas, C., Isobe, M., Russo, G. and Croce, C.M. 1999. Abnormalities at 14q32.1 in T cell malignancies involve two oncogenes. *Proc. Natl. Acad. Sci. USA* 96: 2949-2951.
5. Hallas, C., Pekarsky, Y., Itoyama, T., Varnum, J., Bichi, R., Rothstein, J. and Croce, C.M. 1999. Genomic analysis of human and mouse TCL1 loci reveals a complex of tightly clustered genes. *Proc. Natl. Acad. Sci. USA* 96: 14418-14423.
6. Pekarsky, Y., Koval, A., Hallas, C., Bichi, R., Tresini, M., Malstrom, S., Russo, G., Tsichlis, P. and Croce, C.M. 2000. TCL-1 enhances Akt kinase activity and mediates its nuclear translocation. *Proc. Natl. Acad. Sci. USA* 97: 3028-3033.

CHROMOSOMAL LOCATION

Genetic locus: Tcl1b4 (mouse) mapping to 12 E.

PRODUCT

TCL-1B4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCL-1B4 shRNA Plasmid (m): sc-42995-SH and TCL-1B4 shRNA (m) Lentiviral Particles: sc-42995-V as alternate gene silencing products.

For independent verification of TCL-1B4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42995A, sc-42995B and sc-42995C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TCL-1B4 siRNA (m) is recommended for the inhibition of TCL-1B4 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCL-1B4 gene expression knockdown using RT-PCR Primer: TCL-1B4 (m)-PR: sc-42995-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.