

Ep-CAM siRNA (m): sc-43033

BACKGROUND

The epithelial cell adhesion molecule Ep-CAM, which is also designated tumor-associated calcium signal transducer 1 and MK-1, is a monomeric membrane glycoprotein that is expressed in most normal human epithelium and carcinomas. The human Ep-CAM gene encodes a 314 amino acid protein that is expressed as two forms, a major form and a minor form, which are reduced upon treatment with the amino-glycosylation inhibitor tunicamycin. Ep-CAM is overexpressed in a variety of carcinomas and is, therefore, a potential target for the visualization and therapy of human solid tumors. Ep-CAM contains an extracellular domain containing two epidermal growth factor-like repeats, followed by a cysteine poor region, which is necessary for the adhesion properties of the molecule.

REFERENCES

1. Farr, A., et al. 1991. Epithelial heterogeneity in the murine thymus: a cell surface glycoprotein expressed by subcapsular and medullary epithelium. *J. Histochem. Cytochem.* 39: 645-653.
2. Bergsagel, P.L., et al. 1992. A murine cDNA encodes a pan-epithelial glycoprotein that is also expressed on plasma cells. *J. Immunol.* 148: 590-596.
3. Bjork, P., et al. 1993. Isolation, partial characterization, and molecular cloning of a human colon adenocarcinoma cell-surface glycoprotein recognized by the C215 mouse monoclonal antibody. *J. Biol. Chem.* 268: 24232-24241.
4. Nelson, A.J., et al. 1996. The murine homolog of human Ep-CAM, a homotypic adhesion molecule, is expressed by thymocytes and thymic epithelial cells. *Eur. J. Immunol.* 26: 401-408.
5. Litvinov, S.V., et al. 1997. Epithelial cell adhesion molecule (Ep-CAM) modulates cell-cell interactions mediated by classic cadherins. *J. Cell Biol.* 139: 1337-1348.
6. Taguchi, N., et al. 1999. Abnormal thymic expression of epithelial cell adhesion molecule (Ep-CAM) in New Zealand Black (NZB) mice. *J. Autoimmun.* 13: 393-404.
7. Tomita, Y., et al. 2000. Molecular identification of a human carcinoma-associated glycoprotein antigen recognized by mouse monoclonal antibody FU-MK-1. *Jpn. J. Cancer Res.* 91: 231-238.
8. Trebak, M., et al. 2001. Oligomeric state of the colon carcinoma-associated glycoprotein GA733-2 (Ep-CAM/EGP40) and its role in GA733-mediated homotypic cell-cell adhesion. *J. Biol. Chem.* 276: 2299-2309.
9. Balzar, M., et al. 2001. Epidermal growth factor-like repeats mediate lateral and reciprocal interactions of Ep-CAM molecules in homophilic adhesions. *Mol. Cell. Biol.* 21: 2570-2580.

CHROMOSOMAL LOCATION

Genetic locus: Epcam (mouse) mapping to 17 E4.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Ep-CAM siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ep-CAM shRNA Plasmid (m): sc-43033-SH and Ep-CAM shRNA (m) Lentiviral Particles: sc-43033-V as alternate gene silencing products.

For independent verification of Ep-CAM (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43033A, sc-43033B and sc-43033C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ep-CAM siRNA (m) is recommended for the inhibition of Ep-CAM expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Ep-CAM (G8.8): sc-53532 is recommended as a control antibody for monitoring of Ep-CAM gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Ep-CAM gene expression knockdown using RT-PCR Primer: Ep-CAM (m)-PR: sc-43033-PR (20 μ l, 591 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.