

# β ig-h3 siRNA (m): sc-43124

## BACKGROUND

Human β ig-h3 (α3/β1 Integrin, keratoepithelin) is a secreted, 683 amino acid, transforming growth factor-inducible, extracellular matrix adhesion molecule. β ig-h3 contains an amino-terminal secretory sequence and a carboxy-terminal integrin-binding Arg-Gly-Asp (RGD) domain. β ig-h3 is implicated in mechanisms leading to proliferation, differentiation, wound healing and morphogenesis of corneal tissues. Mutations in the β ig-h3 gene, along with elevated levels of β ig-h3 protein in human corneas, occurs with granular dystrophy (GCD) and other inherited disorders of the cornea. β ig-h3 is also a structural component of the human bladder extracellular matrix and may influence nuclear regulatory or structural functions.

## REFERENCES

1. Skonier, J., Neubauer, M., Madisen, L., Bennett, K., Plowman, G.D. and Purchio, A.F. 1992. cDNA cloning and sequence analysis of β ig-h3, a novel gene induced in a human adenocarcinoma cell line after treatment with transforming growth factor β. *DNA Cell Biol.* 11: 511-522.
2. LeBaron, R.G., Bezverkov, K.I., Zimmer, M.P., Pavelec, R., Skonier, J. and Purchio, A.F. 1995. β ig-h3, a novel secretory protein inducible by transforming growth factor β, is present in normal skin and promotes the adhesion and spreading of dermal fibroblasts *in vitro*. *J. Invest. Dermatol.* 104: 844-849.
3. Rawe, I.M., Zhan, Q., Burrows, R., Bennett, K. and Cintron, C. 1997. β-ig. Molecular cloning and *in situ* hybridization in corneal tissues. *Invest. Ophthalmol. Vis. Sci.* 38: 893-900.
4. Tsujikawa, M., Shimomura, Y., Okada, M., Yamamoto, S., Tano, Y. and Kurahashi, H. 1998. Novel polymorphisms in the β ig-h3 gene. *J. Hum. Genet.* 43: 214-225.
5. Bron, A.J. 2000. Genetics of the corneal dystrophies: what we have learned in the past twenty-five years. *Cornea* 19: 699-711.
6. Billings, P.C., Herrick, D.J., Kucich, U., Engelsberg, B.N., Abrams, W.R., Macarak, E.J., Rosenbloom, J. and Howard, P.S. 2000. Extracellular matrix and nuclear localization of β ig-h3 in human bladder smooth muscle and fibroblast cells. *J. Cell. Biochem.* 79: 261-273.

## CHROMOSOMAL LOCATION

Genetic locus: Tgfb1 (mouse) mapping to 13 B1.

## PRODUCT

β ig-h3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see β ig-h3 shRNA Plasmid (m): sc-43124-SH and β ig-h3 shRNA (m) Lentiviral Particles: sc-43124-V as alternate gene silencing products.

For independent verification of β ig-h3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43124A, sc-43124B and sc-43124C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

β ig-h3 siRNA (m) is recommended for the inhibition of β ig-h3 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor β ig-h3 gene expression knockdown using RT-PCR Primer: β ig-h3 (m)-PR: sc-43124-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.