

Cytokeratin 20 siRNA (h): sc-43313

BACKGROUND

Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue, where they constitute up to 85% of mature keratinocytes in the vertebrate epidermis. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. The α -helical coiled-coil dimers associate laterally end-to-end to form 10 nm diameter filaments. Cytokeratins are useful markers of tissue differentiation, and in addition, they aid in the characterization of malignant tumors. Cytokeratin 20 is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract, and Cytokeratin 20 is a useful marker of pancreatic and colorectal cancer. Cytokeratin 20 is also helpful in distinguishing different types of highly related carcinomas, such as renal oncocytomas from renal cell carcinomas.

REFERENCES

1. van der Velden, L.A., et al. 1993. Cytokeratin expression in normal and (pre)malignant head and neck epithelia: an overview. *Head Neck* 15: 133-146.
2. Moll, R., et al. 1993. The human gene encoding cytokeratin 20 and its expression during fetal development and in gastrointestinal carcinomas. *Differentiation* 53: 75-93.
3. Wauters, C.C., et al. 1995. Keratins 7 and 20 as diagnostic markers of carcinomas metastatic to the ovary. *Hum. Pathol.* 26: 852-855.
4. Marceau, N., et al. 1995. Cytokeratin expression, fibrillar organization and subtle function in liver cells. *Biochem. Cell Biol.* 73: 619-625.
5. Fuchs, E. 1995. Keratins and the skin. *Annu. Rev. Cell Dev. Biol.* 11: 123-153.
6. Quillien, V., et al. 1995. Serum and tissue distribution of a fragment of cytokeratin 19 (cyfra 21-1) in lung cancer patients. *Anticancer Res.* 15: 2857-2863.
7. Mukhopadhyay, T. and Roth, J.A. 1996. Functional inactivation of p53 by antisense RNA induces invasive ability of lung carcinoma cells and down-regulates cytokeratin synthesis. *Anticancer Res.* 16: 1683-1689.

CHROMOSOMAL LOCATION

Genetic locus: KRT20 (human) mapping to 17q21.2.

PRODUCT

Cytokeratin 20 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Cytokeratin 20 shRNA Plasmid (h): sc-43313-SH and Cytokeratin 20 shRNA (h) Lentiviral Particles: sc-43313-V as alternate gene silencing products.

For independent verification of Cytokeratin 20 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43313A, sc-43313B and sc-43313C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Cytokeratin 20 siRNA (h) is recommended for the inhibition of Cytokeratin 20 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Cytokeratin 20 (E-9): sc-271183 is recommended as a control antibody for monitoring of Cytokeratin 20 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG λ BP-HRP: sc-516132 or m-IgG λ BP-HRP (Cruz Marker): sc-516132-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG λ BP-FITC: sc-516185 or m-IgG λ BP-PE: sc-516186 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Cytokeratin 20 gene expression knockdown using RT-PCR Primer: Cytokeratin 20 (h)-PR: sc-43313-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.