

KIF3C siRNA (m): sc-43379

BACKGROUND

The kinesins constitute a large family of microtubule-dependent motor proteins, which are responsible for the distribution of numerous organelles, vesicles and macromolecular complexes throughout the cell. Individual kinesin members play crucial roles in cell division, intracellular transport and membrane trafficking events including endocytosis and transcytosis. Members of the heterotrimeric kinesin II family of microtubule associated motors generally contain two different motor subunits from the KIF3 family, which includes KIF3A, B and C. KIF3 isoforms mediate anterograde transport of membrane bound organelles in neurons and melanosomes, transport between the endoplasmic reticulum and the Golgi, and transport of protein complexes within cilia and flagella required for their morphogenesis. The human KIF3C gene maps to chromosome 2p23.3 and encodes a 793 amino acid protein that is highly expressed in neural tissues such as brain, spinal cord and retina. The selective expression of KIF3C protein in the nervous system during embryonic development and its upregulation during neuroblastoma differentiation suggests a role for this motor during maturation of neuronal cells.

REFERENCES

1. Hamm-Alvarez, S.F. 1998. Molecular motors and their role in membrane traffic. *Adv. Drug Deliv. Rev.* 29: 229-242.
2. Yang, Z. and Goldstein, L.S. 1998. Characterization of the KIF3C neural kinesin-like motor from mouse. *Mol. Biol. Cell* 9: 249-261.
3. Telford, E.A., Wightman, P., Leek, J., Markham, A.F., Lench, N.J. and Bonthon, D.T. 1998. cDNA cloning, genomic organization, and chromosomal localization of a novel human gene that encodes a kinesin-related protein highly similar to mouse KIF3C. *Biochem. Biophys. Res. Commun.* 242: 407-412.
4. Cole, D.G. 1999. Kinesin-II, the heteromeric kinesin. *Cell. Mol. Life Sci.* 56: 217-226.
5. Hirokawa, N. 2000. Stirring up development with the heterotrimeric kinesin KIF3. *Traffic* 1: 29-34.
6. Yang, Z., Xia, C., Roberts, E.A., Bush, K., Nigam, S.K. and Goldstein, L.S. 2001. Molecular cloning and functional analysis of mouse C-terminal kinesin motor KIF3C. *Mol. Cell. Biol.* 21: 765-770.
7. Yang, Z., Roberts, E.A. and Goldstein, L.S. 2001. Functional analysis of mouse kinesin motor KIF3C. *Mol. Cell. Biol.* 21: 5306-5311.
8. Navone, F., Consalez, G.G., Sardella, M., Caspani, E., Pozzoli, O., Frassoni, C., Morlacchi, E., Sitia, R., Sprocati, T. and Cabibbo, A. 2001. Expression of KIF3C kinesin during neural development and *in vitro* neuronal differentiation. *J. Neurochem.* 77: 741-753.
9. Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 603060. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Kif3c (mouse) mapping to 12 A1.1.

PRODUCT

KIF3C siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KIF3C shRNA Plasmid (m): sc-43379-SH and KIF3C shRNA (m) Lentiviral Particles: sc-43379-V as alternate gene silencing products.

For independent verification of KIF3C (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43379A, sc-43379B and sc-43379C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KIF3C siRNA (m) is recommended for the inhibition of KIF3C expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KIF3C gene expression knockdown using RT-PCR Primer: KIF3C (m)-PR: sc-43379-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.