

# $\gamma$ 2-Syntrophin siRNA (m): sc-43444

## BACKGROUND

The syntrophins are structurally related PDZ-domain-containing proteins that facilitate the recruitment of signaling proteins, such as NOS1, to the cell membrane. Syntrophins associate directly with dystrophin, a scaffold protein that is part of a complex which is disrupted in muscular dystrophy, and with dystrophin-related proteins.  $\gamma$ 2-Syntrophin, also known as SYN5 or G2SYN, is a 539 amino acid member of the syntrophin family that functions as an adaptor protein to link and organize various proteins, such as dystrophin, within the cell.  $\gamma$ 2-Syntrophin, which is localized to the sarcolemma cell membrane, is widely expressed and contains one PH domain and one PDZ domain.  $\gamma$ 2-Syntrophin is able to recruit proteins to the membrane through its PDZ domain, which is unavailable when the protein is bound to a substrate. As a result of its interaction with various proteins,  $\gamma$ 2-Syntrophin is implicated in inherited muscular dystrophy and in the development of autism.

## REFERENCES

1. Newey, S.E., Benson, M.A., Ponting, C.P., Davies, K.E. and Blake, D.J. 2000. Alternative splicing of dystrobrevin regulates the stoichiometry of syntrophin binding to the dystrophin protein complex. *Curr. Biol.* 10: 1295-1298.
2. Abdelmoity, A., Padre, R.C., Burzynski, K.E., Stull, J.T. and Lau, K.S. 2000. Neuronal nitric oxide synthase localizes through multiple structural motifs to the sarcolemma in mouse myotubes. *FEBS Lett.* 482: 65-70.
3. Adams, M.E., Kramarcy, N., Krall, S.P., Rossi, S.G., Rotundo, R.L., Sealock, R. and Froehner, S.C. 2000. Absence of  $\alpha$ -Syntrophin leads to structurally aberrant neuromuscular synapses deficient in utrophin. *J. Cell Biol.* 150: 1385-1398.
4. Ort, T., Maksimova, E., Dirx, R., Kachinsky, A.M., Berghs, S., Froehner, S.C. and Solimena, M. 2000. The receptor tyrosine phosphatase-like protein ICA512 binds the PDZ domains of  $\beta$ 2-Syntrophin and nNOS in pancreatic  $\beta$  cells. *Eur. J. Cell Biol.* 79: 621-630.
5. Rocco, P., Vainzof, M., Froehner, S.C., Peters, M.F., Marie, S.K., Passos-Bueno, M.R. and Zatz, M. 2000. Brazilian family with pure autosomal dominant spastic paraplegia maps to 8q: analysis of muscle  $\beta$ 1-Syntrophin. *Am. J. Med. Genet.* 92: 122-127.

## CHROMOSOMAL LOCATION

Genetic locus: Sntg2 (mouse) mapping to 12 B1.

## PRODUCT

$\gamma$ 2-Syntrophin siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see  $\gamma$ 2-Syntrophin shRNA Plasmid (m): sc-43444-SH and  $\gamma$ 2-Syntrophin shRNA (m) Lentiviral Particles: sc-43444-V as alternate gene silencing products.

For independent verification of  $\gamma$ 2-Syntrophin (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43444A, sc-43444B and sc-43444C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

$\gamma$ 2-Syntrophin siRNA (m) is recommended for the inhibition of  $\gamma$ 2-Syntrophin expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor  $\gamma$ 2-Syntrophin gene expression knockdown using RT-PCR Primer:  $\gamma$ 2-Syntrophin (m)-PR: sc-43444-PR (20  $\mu$ l, 440 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.