

TCP-1 ζ 2 siRNA (m): sc-43456

BACKGROUND

The protein TCP-1 (t complex polypeptide 1) is a subunit of the hetero-oligomeric complex CCT (chaperonin containing TCP-1) present in the eukaryotic cytosol. The CCT of eukaryotic cytosol is composed of eight different subunit species, TCP-1 α , β , γ , δ , ϵ , ζ , η and θ , each encoded by a different gene. Two ζ subunits have been described: TCP-1 ζ (also designated TCP-1 ζ 1) and TCP-1 ζ 2. TCP-1 subunits are proposed to have independent functions in folding its *in vivo* substrates, the actins and tubulins. TCP-1 was first identified in the mouse as relevant for tail-less and embryonic lethal phenotypes. Sequences homologous to TCP-1 have been isolated in several other species, and the yeast TCP-1 has been shown to encode a molecular chaperone for Actin and Tubulin. TCP-1 found in mammalian cells and yeast plays an important role in the folding of cytosolic proteins.

REFERENCES

1. Ahnert, V., May, C., Gerke, R. and Kindl, H. 1996. Cucumber T complex protein. Molecular cloning, bacterial expression and characterization within a 22-S cytosolic complex in cotyledons and hypocotyls. *Eur. J. Biochem.* 235: 114-119.
2. Iijima, M., Shimizu, H., Tanaka, Y. and Urushihara, H. 1998. A Dictyostelium discoideum homologue to TCP-1 is essential for growth and development. *Gene* 213: 101-106.
3. Ritco-Vonsovici, M. and Willison, K.R. 2000. Defining the eukaryotic cytosolic chaperonin-binding sites in human tubulins. *J. Mol. Biol.* 304: 81-98.
4. Hynes, G.M. and Willison, K.R. 2000. Individual subunits of the eukaryotic cytosolic chaperonin mediate interactions with binding sites located on subdomains of β -Actin. *J. Biol. Chem.* 275: 18985-18994.
5. Campos, E.G. and Hamdan, F.F. 2000. Cloning of the chaperonin T complex polypeptide 1 gene from *Schistosoma mansoni* and studies of its expression levels under heat shock and oxidative stress. *Parasitol. Res.* 86: 253-258.
6. Yokota, S.I., Yanagi, H., Yura, T. and Kubota, H. 2000. Upregulation of cytosolic chaperonin CCT subunits during recovery from chemical stress that causes accumulation of unfolded proteins. *Eur. J. Biochem.* 267: 1658-1664.

CHROMOSOMAL LOCATION

Genetic locus: Cct6b (mouse) mapping to 11 C.

PRODUCT

TCP-1 ζ 2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCP-1 ζ 2 shRNA Plasmid (m): sc-43456-SH and TCP-1 ζ 2 shRNA (m) Lentiviral Particles: sc-43456-V as alternate gene silencing products.

For independent verification of TCP-1 ζ 2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43456A, sc-43456B and sc-43456C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TCP-1 ζ 2 siRNA (m) is recommended for the inhibition of TCP-1 ζ 2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCP-1 ζ 2 gene expression knockdown using RT-PCR Primer: TCP-1 ζ 2 (m)-PR: sc-43456-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.