

# Taq DNA Polymerase: sc-43501

## PRODUCT DESCRIPTION

*Taq* DNA Polymerase is a highly thermostable DNA polymerase of the thermophilic bacterium *Thermus aquaticus*. The enzyme catalyzes 5'→3' synthesis of DNA, has no detectable 3'→5' exonuclease (proofreading) activity, and possesses 5'→3' exonuclease activity. In addition, *Taq* DNA Polymerase exhibits deoxynucleotidyl transferase activity, which frequently results in the addition of extra adenines at the 3'-end of PCR products. Recombinant *Taq* DNA Polymerase is ideal for standard PCR of amplicons 5 kb or shorter.

COMPONENT	QUANTITY
<i>Taq</i> DNA Polymerase	500 U
10X <i>Taq</i> Buffer with KCl	2 x 1.25 ml
10X <i>Taq</i> Buffer with (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	2 x 1.25 ml
25 mM MgCl <sub>2</sub> *	2 x 1.25 ml

### APPLICATIONS

- Routine PCR amplification of DNA fragments up to 5 kb
- Generation of PCR product for TA cloning
- DNA labeling
- DNA sequencing

### SOURCE

*Taq* DNA Polymerase is derived from *E. coli* cells with a cloned *pol* gene from *Thermus aquaticus* YT1.

### DEFINITION OF ACTIVITY UNIT

One unit of the *Taq* DNA Polymerase enzyme catalyzes the incorporation of 10 nmol of deoxyribonucleotides into a polynucleotide fraction (adsorbed on DE-81) in 30 minutes at 70° C.

Enzyme activity is assayed in the following mixture: 67 mM Tris-HCl (pH 8.8 at 25° C), 6.7 mM MgCl<sub>2</sub>, 1 mM 2-mercaptoethanol, 50 mM NaCl, 0.1 mg/ml BSA, 0.75 mM activated salmon milt DNA, 0.2 mM of each dNTP, 0.4 MBq/ml [<sup>3</sup>H]-dTTP.

### STORAGE BUFFER

The enzyme is supplied in 20 mM Tris-HCl (pH 8.0), 1 mM DTT, 0.1 mM EDTA, 100 mM KCl, 0.5% (v/v) Nonidet<sup>®</sup> P40, 0.5% (v/v) Tween<sup>®</sup> 20 and 50% (v/v) glycerol.

### 10X *Taq* BUFFER WITH KCl

100 mM Tris-HCl (pH 8.8 at 25° C), 500 mM KCl, 0.8% (v/v) Nonidet P40.

### 10X *Taq* BUFFER WITH (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>

750 mM Tris-HCl (pH 8.8 at 25° C), 200 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.1% (v/v) Tween 20.

### INHIBITION AND INACTIVATION

Inhibited by ionic detergents (deoxycholate, sarkosyl and SDS) at concentrations higher than 0.06, 0.02 and 0.01%, respectively.

Inactivated by phenol/chloroform extraction.

## PROTOCOL

To prepare several parallel reactions and to minimize the possibility of pipetting errors, prepare a PCR master mix by mixing water, buffer, dNTPs, primers and *Taq* DNA Polymerase. Prepare sufficient master mix for the number of reactions plus one extra. Aliquot the master mix into individual PCR tubes and then add template DNA.

- Gently vortex and briefly centrifuge all solutions after thawing.
- Place a thin-walled PCR tube on ice and add the following components for each 50 µl reaction:

10X <i>Taq</i> Buffer	5 µl
dNTP Mix, 2 mM each	5 µl (0.2 mM of each)
Forward primer	0.1-1.0 µM
Reverse primer	0.1-1.0 µM
25 mM MgCl <sub>2</sub> *	1-4 mM
Template DNA	10 pg-1 µg
<i>Taq</i> DNA Polymerase	1.25 U
Water, nuclease-free	to 50 µl
Total volume	50 µl

\*Volumes of 25 mM MgCl<sub>2</sub>, required for specific final MgCl<sub>2</sub> concentration:

Final concentration of MgCl <sub>2</sub> , mM	1	1.5	2	2.5	3	4
Volume of 25 mM MgCl <sub>2</sub> to be added for 50 µl reaction, µl	2	3	4	5	6	8

- Gently vortex the samples and spin down.
- If using a thermal cycler that does not use a heated lid, overlay the reaction mixture with 25 µL of mineral oil.
- Perform PCR using recommended thermal cycling conditions:

STEP	TEMP° C	TIME	NO. OF CYCLES
Initial denaturation	95	1-3 min.	1
Denaturation	95	30 sec.	25-40
Annealing	T <sub>m</sub> -5	30 sec.	25-40
Extension	72	1 min/kb	25-40
Final Extension	72	5-15 min	1

## PREVENTING CONTAMINATION OF PCR REACTION

During PCR more than 10 million copies of template DNA are generated. Therefore, care must be taken to avoid contamination with other templates and amplicons that may be present in the laboratory environment. General recommendations to lower the risk of contamination are as follows:

- Prepare your DNA sample, set up the PCR mixture, perform thermal cycling and analyze PCR products in separate areas.
- Set up PCR mixtures in a laminar flow cabinet equipped with an UV lamp.
- Wear fresh gloves for DNA purification and reaction set up.
- Use reagent containers dedicated for PCR. Use positive displacement pipettes, or use pipette tips with aerosol filters to prepare DNA samples and perform PCR set up.
- Use PCR-certified reagents, including high quality water.
- Always perform “no template control” (NTC) reactions to check for contamination.

## GUIDELINES FOR PRIMER DESIGN

General recommendations for PCR primer design as outlined below:

- PCR primers are generally 15-30 nucleotides long.
- Optimal GC content of the primer is 40-60%. Ideally, C and G nucleotides should be distributed uniformly along the primer.
- Avoid placing more than three G or C nucleotides at the 3'-end to lower the risk of non-specific priming. If possible, the primer should terminate with a G or C at the 3'-end.
- Avoid self-complementary primer regions, complementarities between the primers and direct primer repeats to prevent hairpin formation and primer dimerization.
- Check for possible sites of undesired complementary between primers and template DNA.
- When designing degenerate primers, place at least 3 conserved nucleotides at the 3'-end.
- Differences in melting temperatures ( $T_m$ ) between the two primers should not exceed 5° C.

### ESTIMATION OF PRIMER MELTING TEMPERATURE

For primers containing less than 25 nucleotides, the approximate melting temperature ( $T_m$ ) can be calculated using the following equation:

$$T_m = 4(G + C) + 2(A + T)$$

where G, C, A, T represent the number of respective nucleotides in the primer.

If the primer contains more than 25 nucleotides specialized computer programs are recommended to account for interactions of adjacent bases, effect of salt concentration, etc.

## COMPONENTS OF THE REACTION MIXTURE

### TEMPLATE DNA

Optimal amounts of template DNA in the 50  $\mu$ l reaction volume are 0.01 ng-1 ng for both plasmid and phage DNA, and 0.1  $\mu$ g-1  $\mu$ g for genomic DNA. Higher amount of template increases the risk of generation of non-specific PCR products. Lower amount of template reduces the accuracy of the amplification.

All routine DNA purification methods are suitable for template preparation. Trace amounts of certain agents used for DNA purification, such as phenol, EDTA and proteinase K, can inhibit DNA polymerases. Ethanol precipitation and repeated washes of the DNA pellet with 70% ethanol normally removes trace contaminants from DNA samples.

### MgCl<sub>2</sub> CONCENTRATION

Due to the binding of Mg<sup>2+</sup> to dNTPs, primers and DNA templates, Mg<sup>2+</sup> concentration needs to be optimized for maximal PCR yield. The recommended concentration range is 1-4 mM. If the Mg<sup>2+</sup> concentration is too low, the

yield of PCR product could be reduced. On the contrary, non-specific PCR products may appear and the PCR fidelity may be reduced if the Mg<sup>2+</sup> concentration is too high.

If the DNA samples contain EDTA or other metal chelators, the Mg<sup>2+</sup> ion concentration in the PCR mixture should be increased accordingly (one molecule of EDTA binds one Mg<sup>2+</sup>).

### dNTPS

The recommended final concentration of each dNTP is 0.2 mM. In certain PCR applications, higher dNTP concentrations may be necessary. Due to the binding of Mg<sup>2+</sup> to dNTPs, the MgCl<sub>2</sub> concentration needs to be adjusted accordingly. It is essential to have equal concentrations of all four nucleotides (dATP, dCTP, dGTP and dTTP) present in the reaction mixture.

To achieve 0.2 mM concentration of each dNTP in the PCR mixture, use the following volumes of dNTP mixes:

VOLUME OF PCR MIXTURE	dNTP MIX, 2 mM EACH	dNTP MIX, 10 mM EACH	dNTP MIX, 25 mM EACH
50 $\mu$ l	5 $\mu$ l	1 $\mu$ l	0.4 $\mu$ l
25 $\mu$ l	2.5 $\mu$ l	0.5 $\mu$ l	0.2 $\mu$ l
20 $\mu$ l	2 $\mu$ l	0.5 $\mu$ l	0.16 $\mu$ l

### PRIMERS

The recommended concentration range of the PCR primers is 0.1  $\mu$ M-1  $\mu$ M. Excessive primer concentrations increase the probability of mispriming and generation of non-specific PCR products.

For degenerate primers, primer concentrations in the range of 0.3  $\mu$ M-1  $\mu$ M are often favorable.

## CYCLING PARAMETERS

### INITIAL DNA DENATURATION

It is essential to completely denature the template DNA at the beginning of PCR to ensure efficient utilization of the template during the first amplification cycle. If the GC content of the template is 50% or less, an initial 193 minute denaturation at 95° C is sufficient. For GC9-rich templates this step should be prolonged up to 10 minutes. If longer initial denaturation is required, or the DNA is denatured at a higher temperature, *Taq* DNA Polymerase should be added after the initial denaturation step to avoid a decrease in its activity.

### DENATURATION

A DNA denaturation time of 30 seconds per cycle at 95° C is normally sufficient. For GC9-rich DNA templates, this step can be prolonged to 394 minutes. DNA denaturation can also be enhanced by the addition of either 10-15% glycerol or 10% DMSO, 5% formamide or 1-1.5 M betaine. The melting temperature of the primer-template complex decreases significantly in the presence of these reagents. Therefore, the annealing temperature has to be adjusted accordingly. In addition, 10% DMSO and 5% formamide inhibit DNA polymerases by 50%. Thus, the amount of the enzyme should be increased if these additives are used.

### PRIMER ANNEALING

The annealing temperature should be 5° C lower than the melting temperature ( $T_m$ ) of the primers. Annealing for 30 seconds is normally sufficient. If non-specific PCR products appear, the annealing temperature should be optimized stepwise in 192° C increments. When additives that will change the melting temperature of the primer-template complex are used (glycerol, DMSO, formamide and betaine), the annealing temperature must also be adjusted.

#### EXTENSION

The optimal extension temperature for *Taq* DNA Polymerase is 70-75° C. The recommended extension step is 1 minute at 72° C for PCR products up to 2 kb. For larger products, the extension time should be prolonged by 1 minute/kb.

#### NUMBER OF CYCLES

The number of cycles may vary depending on the amount of template DNA in the PCR mixture and the expected PCR product yield.

If less than 10 copies of the template are present in the reaction, about 40 cycles are required. For higher template amounts, 25-35 cycles are sufficient.

#### FINAL EXTENSION

After the last cycle, it is recommended to incubate the PCR mixture at 72° C for an additional 5-15 minutes to fill-in any possible incomplete reaction products. If the PCR product will be cloned into TA vectors, the final extension step may be prolonged to 30 minutes to ensure the highest efficiency of 3'-dA tailing of PCR product.

### ADDITIONAL PRODUCT INFORMATION

#### NOTICE TO PURCHASER

Use of this product is covered by US Patent No. 6,127,155. The purchase of this product includes a limited, non-transferable immunity from suit under the foregoing patent claims for using only this amount of product for the purchaser's own internal research. No right under any other patent claim, no right to perform any patented method and no right to perform commercial services of any kind, including without limitation reporting the results of purchaser's activities for a fee or other commercial consideration, is conveyed expressly, by implication, or by estoppel.

This product is for research use only. Diagnostic uses under Roche patents require a separate license from Roche. Further information on purchasing licenses may be obtained by contacting:

Director of Licensing  
Applied Biosystems  
850 Lincoln Centre Drive  
Foster City, CA 94404-1128

#### PRODUCT USE LIMITATION

This product is developed, designed and sold exclusively for research purposes and *in vitro* use only. The product was not tested for use in diagnostics or for drug development, nor is it suitable for administration to humans or animals.