

Pontin 52 siRNA (h): sc-43543

BACKGROUND

Pontin 52 is a nuclear matrix protein that is primarily expressed in the nucleus and is also present in the cytoplasm. Pontin 52 is expressed in the nucleoplasm of whole cells, but is not present in the nucleoli. Pontin 52 is also designated RUVBL1, for *E. coli* RuvB-like 1 protein, or NMP 238, and is the human homolog of rat TIP49. Pontin 52 contains an ATPase/helicase motif and may represent a class of cofactors recruited by transcriptional activation domains that function in diverse pathways. For instance, *in vivo*, Pontin 52 is complexed with Myc and Reptin 52, which is a Pontin 52 related protein also designated RUVBL2. The interaction of Pontin 52 with Myc is dependent upon a Myc domain essential for oncogenic activity, suggesting that functional Pontin 52 is an essential mediator of Myc oncogenic transformation. The gene encoding human Pontin 52 maps to chromosome 3q21.3.

REFERENCES

1. Bauer, A., et al. 1998. Pontin 52, an interaction partner of β -catenin, binds to the TATA box-binding protein. *Proc. Natl. Acad. Sci. USA* 95: 14787-14792.
2. Makino, Y., et al. 1998. TIP49, homologous to the bacterial DNA helicase RuvB, acts as an autoantigen in human. *Biochem. Biophys. Res. Commun.* 245: 819-823.
3. Holzmann, K., et al. 1998. Identification and characterization of the ubiquitously occurring nuclear matrix protein NMP 238. *Biochem. Biophys. Res. Commun.* 252: 39-45.
4. Qiu, X.B., et al. 1998. An eukaryotic RuvB-like protein (RUVBL1) essential for growth. *J. Biol. Chem.* 273: 27786-27793.

CHROMOSOMAL LOCATION

Genetic locus: RUVBL1 (human) mapping to 3q21.3.

PRODUCT

Pontin 52 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Pontin 52 shRNA Plasmid (h): sc-43543-SH and Pontin 52 shRNA (h) Lentiviral Particles: sc-43543-V as alternate gene silencing products.

For independent verification of Pontin 52 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43543A, sc-43543B and sc-43543C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Pontin 52 siRNA (h) is recommended for the inhibition of Pontin 52 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Pontin 52 (A-11): sc-393905 is recommended as a control antibody for monitoring of Pontin 52 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Pontin 52 gene expression knockdown using RT-PCR Primer: Pontin 52 (h)-PR: sc-43543-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Park, E.J., et al. 2011. The human Ino80 binds to microtubule via the E-hook of Tubulin: implications for the role in spindle assembly. *Biochem. Biophys. Res. Commun.* 416: 416-420.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.