

Calponin 1/2/3 siRNA (h): sc-43657

BACKGROUND

Calponin regulates smooth muscle cell contraction and is a marker of smooth muscle cell differentiation. Calponin, an Actin- and Tropomyosin-binding protein, is characterized as an inhibitory factor of smooth-muscle actomyosin activity. Calponin is implicated in the regulation of smooth muscle contraction through its interaction with F-Actin and inhibition of the Actin-activated MgATPase activity of phosphorylated myosin. Both properties are lost following phosphorylation (primarily at Serine 175) by protein kinase C or calmodulin-dependent protein kinase II. The three forms of Calponin, Calponin 1 (basic Calponin), Calponin 2 (neutral Calponin) and Calponin 3 (acidic Calponin) are found in smooth muscle tissue. Additionally, Calponin 2 is found in heart muscle tissue and Calponin 3 is found in the brain.

REFERENCES

1. Tang, D.C., et al. 1996. Structure-function relations of smooth muscle Calponin. The critical role of Serine 175. *J. Biol. Chem.* 271: 8605-8611.
2. Masuda, H., et al. 1996. Molecular cloning and characterization of human non-smooth muscle Calponin. *J. Biochem.* 120: 415-424.
3. Doi, M., et al. 1997. Reduced expression of Calponin in canine basilar artery after subarachnoid haemorrhage. *Acta Neurochir.* 139: 77-81.
4. Kaneko, T., et al. 2000. Identification of Calponin as a novel substrate of Rho-kinase. *Biochem. Biophys. Res. Commun.* 273: 110-116.
5. di Gioia, C.R., et al. 2000. Angiotensin II increases Calponin expression in cultured rat vascular smooth muscle cells. *Biochem. Biophys. Res. Commun.* 279: 965-969.
6. Yoshimoto, R., et al. 2000. Proteolysis of acidic Calponin by μ -calpain. *J. Biochem.* 128: 1045-1049.

PRODUCT

Calponin 1/2/3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Calponin 1/2/3 shRNA Plasmid (h): sc-43657-SH and Calponin 1/2/3 shRNA (h) Lentiviral Particles: sc-43657-V as alternate gene silencing products.

For independent verification of Calponin 1/2/3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43657A, sc-43657B and sc-43657C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Calponin 1/2/3 siRNA (h) is recommended for the inhibition of Calponin 1/2/3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Calponin 1/2/3 (G-10): sc-136987 is recommended as a control antibody for monitoring of Calponin 1/2/3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.