



# K-Ras siRNA (h2): sc-43874

## BACKGROUND

The mammalian Ras (also designated v-Ha-Ras, Harvey rat sarcoma viral oncogene homolog, HRAS1, K-Ras, N-Ras, RASH1 or c-bas/has) gene family consists of the Harvey and Kirsten Ras genes (c-H-Ras1 and c-K-Ras2), an inactive pseudogene of each (c-H-Ras2 and c-K-Ras1) and the N-Ras gene. The three Ras oncogenes, H-Ras, K-Ras and N-Ras, encode proteins with GTP/GDP binding and GTPase activity. Ras proteins alternate between an inactive form bound to GDP and an active form bound to GTP, activated by a guanine nucleotide-exchange factor (GEF) and inactivated by a GTPase-activating protein (GAP). Ras nomenclature originates from the characterization of human DNA sequences homologous to cloned DNA fragments containing oncogenic sequences of a type C mammalian retrovirus, the Harvey strain of murine sarcoma virus (HaMSV), derived from the rat. Under normal conditions, Ras family members influence cell growth and differentiation events in a subcellular membrane compartmentalization-based signaling system. Oncogenic Ras can deregulate processes that control both cell proliferation and apoptosis. The Ras superfamily of GTP hydrolysis-coupled signal transduction relay proteins can be subclassified into Ras, Rho, Rab and ARF families.

## CHROMOSOMAL LOCATION

Genetic locus: KRAS (human) mapping to 12p12.1.

## PRODUCT

K-Ras siRNA (h2) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see K-Ras shRNA Plasmid (h2): sc-43874-SH and K-Ras shRNA (h2) Lentiviral Particles: sc-43874-V as alternate gene silencing products.

For independent verification of K-Ras (h2) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43874A, sc-43874B and sc-43874C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

K-Ras siRNA (h2) is recommended for the inhibition of K-Ras expression in human cells.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

K-Ras (F234): sc-30 is recommended as a control antibody for monitoring of K-Ras gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor K-Ras gene expression knockdown using RT-PCR Primer: K-Ras (h2)-PR: sc-43874-PR (20  $\mu$ l, 588 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

- Ryu, S.H., et al. 2010. Oncogenic Ras-mediated downregulation of Clast1/LR8 is involved in Ras-mediated neoplastic transformation and tumorigenesis in NIH3T3 cells. *Cancer Sci.* 101: 1990-1996.
- Calvert, R.J., et al. 2013. K-Ras 4A and 4B mRNA levels correlate with superoxide in lung adenocarcinoma cells, while at the protein level, only mutant K-Ras 4A protein correlates with superoxide. *Lung Cancer* 80: 263-269.
- Liu, H., et al. 2019. KRAS-enhanced macropinocytosis and reduced FcRn-mediated recycling sensitize pancreatic cancer to albumin-conjugated drugs. *J. Control. Release* 296: 40-53.
- Shin, D.H., et al. 2023. Oncogenic KRAS mutation confers chemoresistance by upregulating SIRT1 in non-small cell lung cancer. *Exp. Mol. Med.* 55: 2220-2237.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.