

# Mfn2 siRNA (h): sc-43928

## BACKGROUND

Mitofusin 1 (Mfn1) and mitofusin 2 (Mfn2) are homologs for the *Drosophila* protein fuzzy onion (Fzo). They are mitochondrial membrane proteins and are mediators of mitochondrial fusion. A GTPase domain is required for Mfn protein function but the molecular mechanisms of the GTPase-dependent reaction as well as the functional division of the two Mfn proteins are unknown. They are essential for embryonic development and may play a role in the pathobiology of obesity. Although the Mfn1 and Mfn2 genes are broadly expressed, they show different levels of expression in different tissues. Two Mfn1 transcripts are elevated in heart, while Mfn2 mRNA is abundantly expressed in heart and muscle tissue but present only at low levels in many other tissues. Mfn1 localizes to mitochondria and participates in at least two different high molecular weight protein complexes in a GTP-dependent manner. Purified recombinant Mfn1 exhibited approximately eightfold higher GTPase activity than Mfn2.

## CHROMOSOMAL LOCATION

Genetic locus: MFN2 (human) mapping to 1p36.22.

## PRODUCT

Mfn2 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Mfn2 shRNA Plasmid (h): sc-43928-SH and Mfn2 shRNA (h) Lentiviral Particles: sc-43928-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Mfn2 siRNA (h) is recommended for the inhibition of Mfn2 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

Mfn2 (F-5): sc-515647 is recommended as a control antibody for monitoring of Mfn2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Mfn2 gene expression knockdown using RT-PCR Primer: Mfn2 (h)-PR: sc-43928-PR (20  $\mu$ l, 557 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

- Caino, M.C., et al. 2015. PI3K therapy reprograms mitochondrial trafficking to fuel tumor cell invasion. *Proc. Natl. Acad. Sci. USA* 112: 8638-8643.
- Cheng, C.T., et al. 2016. Metabolic stress-induced phosphorylation of KAP1 Ser473 blocks mitochondrial fusion in breast cancer cells. *Cancer Res.* 76: 5006-5018.
- Corazao-Rozas, P., et al. 2016. Mitochondrial oxidative phosphorylation controls cancer cell's life and death decisions upon exposure to MAPK inhibitors. *Oncotarget* 7: 39473-39485.
- Caino, M.C., et al. 2016. A neuronal network of mitochondrial dynamics regulates metastasis. *Nat. Commun.* 7: 13730.
- Zhang, R., et al. 2017. Upregulation of miR-195 accelerates oxidative stress-induced retinal endothelial cell injury by targeting mitofusin 2 in diabetic rats. *Mol. Cell. Endocrinol.* 452: 33-43.
- Wang, Q., et al. 2017. Deletion of PRKAA triggers mitochondrial fission by inhibiting the autophagy-dependent degradation of DNM1L. *Autophagy* 13: 404-422.
- Lin, H.Y., et al. 2018. The causal role of mitochondrial dynamics in regulating Insulin resistance in diabetes: link through mitochondrial reactive oxygen species. *Oxid. Med. Cell. Longev.* 2018: 7514383.
- Li, X., et al. 2019. FGF21 mediates mesenchymal stem cell senescence via regulation of mitochondrial dynamics. *Oxid. Med. Cell. Longev.* 2019: 4915149.
- He, H., et al. 2019. Vascular progenitor cell senescence in patients with Marfan syndrome. *J. Cell. Mol. Med.* 23: 4139-4152.
- Li, H., et al. 2019. Mst1 deletion attenuates renal ischaemia-reperfusion injury: the role of microtubule cytoskeleton dynamics, mitochondrial fission and the GSK3 $\beta$ -p53 signalling pathway. *Redox Biol.* 20: 261-274.
- Braganza, A., et al. 2019. Myoglobin induces mitochondrial fusion, thereby inhibiting breast cancer cell proliferation. *J. Biol. Chem.* 294: 7269-7282.
- Chang, Y.H., et al. 2020. The causal role of mitochondrial dynamics in regulating innate immunity in diabetes. *Front. Endocrinol.* 11: 445.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.