

MICA/B siRNA (h): sc-43931

BACKGROUND

MICA and MICB are stress-induced antigens that are related to major histocompatibility complex (MHC) class I molecules. MICA and MICB are frequently expressed in epithelial tumors. These highly glycosylated cell surface proteins are stably expressed without conventional class I peptide ligands or association with β -2-Microglobulin. The expression is induced on proliferating or heat shock stressed epithelial cells. MICA and MICB are broadly recognized by intestinal epithelial V δ 1 $\gamma\delta$ T cells expressing variable TCRs, suggesting that these antigens may play a central role in the signaling of cellular distress to evoke immune responses in the intestinal epithelium.

REFERENCES

1. Bahram, S., et al. 1994. A second lineage of mammalian major histocompatibility complex class I genes. *Proc. Natl. Acad. Sci. USA* 91: 6259-6263.
2. Bahram, S., et al. 1996. Nucleotide sequence of the human MHC class I MICA gene. *Immunogenetics* 44: 80-81.
3. Bahram, S., et al. 1996. Genomic structure of the human MHC class I MICB gene. *Immunogenetics* 45: 161-162.
4. Groh, V., et al. 1996. Cell stress-regulated human major histocompatibility complex class I gene expressed in gastrointestinal epithelium. *Proc. Natl. Acad. Sci. USA* 93: 12445-12450.
5. Groh, V., et al. 1998. Recognition of stress-induced MHC molecules by intestinal epithelial $\gamma\delta$ T cells. *Science* 279: 1737-1740.
6. Steinle, A., et al. 1998. Diversification, expression and $\gamma\delta$ T cell recognition of evolutionarily distant members of the MIC family of major histocompatibility complex class I-related molecules. *Proc. Natl. Acad. Sci. USA* 95: 12510-12515.

CHROMOSOMAL LOCATION

Genetic locus: MICA/MICB (human) mapping to 6p21.33.

PRODUCT

MICA/B siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MICA/B shRNA Plasmid (h): sc-43931-SH and MICA/B shRNA (h) Lentiviral Particles: sc-43931-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MICA/B siRNA (h) is recommended for the inhibition of MICA/B expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

MICA/B (F-6): sc-137242 is recommended as a control antibody for monitoring of MICA/B gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MICA/B gene expression knockdown using RT-PCR Primer: MICA/B (h)-PR: sc-43931-PR (20 μ l, 425 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Huang, B., et al. 2011. Modulation of NKG2D-ligand cell surface expression enhances immune cell therapy of cancer. *J. Immunother.* 34: 289-296.
2. Shimizu, T., et al. 2018. Low dose gemcitabine increases the cytotoxicity of human V γ 9V δ 2 T cells in bladder cancer cells *in vitro* and in an orthotopic xenograft model. *Oncoimmunology* 7: e1424671.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.