

# TGFβ1/2/3 siRNA (h): sc-44146

## BACKGROUND

Transforming growth factor βs (TGFβs) were originally discovered due to their ability to promote anchorage-independent growth of rat NRK fibroblasts in the presence of TGFα. It is now realized that TGFβs mediate many cell-cell interactions that occur during embryonic development. Three TGFβs have been identified in mammals. TGFβ1, TGFβ2 and TGFβ3 are each synthesized as precursor proteins that are very similar in that each is cleaved to yield a 112 amino acid polypeptide that remains associated with the latent portion of the molecules. Biologically active TGFβ requires dimerization of the monomers (usually homodimers) and release of the latent peptide portion. Overall, the mature region of the TGFβ3 protein has approximately 80% identity to the mature region of both TGFβ1 and TGFβ2. However, the NH<sub>2</sub> terminals or precursor regions of their molecules share only 27% sequence identity.

## PRODUCT

TGFβ1/2/3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TGFβ1/2/3 shRNA Plasmid (h): sc-44146-SH and TGFβ1/2/3 shRNA (h) Lentiviral Particles: sc-44146-V as alternate gene silencing products.

For independent verification of TGFβ1/2/3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44146A, sc-44146B and sc-44146C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

TGFβ1/2/3 siRNA (h) is recommended for the inhibition of TGFβ1/2/3 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## GENE EXPRESSION MONITORING

TGFβ3 (G-9): sc-166833 is recommended as a control antibody for monitoring of TGFβ1/2/3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

## SELECT PRODUCT CITATIONS

1. Yamauchi, Y., et al. 2013. Lewis lung carcinoma progression is facilitated by TIG-3 fibroblast cells. *Anticancer Res.* 33: 3791-3798.
2. Nagahara, T., et al. 2015. Hepatic stellate cells promote upregulation of epithelial cell adhesion molecule and epithelial-mesenchymal transition in hepatic cancer cells. *Oncol. Rep.* 34: 1169-1177.
3. Chang, T.P., et al. 2015. Bortezomib inhibits expression of TGF-β1, IL-10, and CXCR4, resulting in decreased survival and migration of cutaneous T cell lymphoma cells. *J. Immunol.* 194: 2942-2953.
4. Polimeni, M., et al. 2016. Multi-walled carbon nanotubes directly induce epithelial-mesenchymal transition in human bronchial epithelial cells via the TGF-β-mediated Akt/GSK-3β/SNAIL-1 signalling pathway. *Part. Fibre Toxicol.* 13: 27.
5. Qin, G., et al. 2016. Reciprocal activation between MMP-8 and TGF-β1 stimulates EMT and malignant progression of hepatocellular carcinoma. *Cancer Lett.* 374: 85-95.
6. Sakamoto, A., et al. 2018. Cross-talk between the transcription factor Sp1 and C/EBPβ modulates TGFβ1 production to negatively regulate the expression of chemokine RANTES. *Heliyon* 4: e00679.
7. He, H., et al. 2019. Vascular progenitor cell senescence in patients with Marfan syndrome. *J. Cell. Mol. Med.* 23: 4139-4152.
8. You, W., et al. 2019. TGF-β mediates aortic smooth muscle cell senescence in Marfan syndrome. *Aging* 11: 3574-3584.
9. Yamaguchi, R., et al. 2020. TRIM28/TIF1β and Fli-1 negatively regulate peroxynitrite generation via DUOX2 to decrease the shedding of membrane-bound fractalkine in human macrophages after exposure to substance P. *Cytokine* 134: 155180.
10. Yamaguchi, R., et al. 2022. IL-23 production in human macrophages is regulated negatively by tumor necrosis factor α-induced protein 3 and positively by specificity protein 1 after stimulation of the toll-like receptor 7/8 signaling pathway. *Heliyon* 8: e08887.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.