



Heme Oxygenase 1 siRNA (h2): sc-44306

BACKGROUND

Heme oxygenases are microsomal enzymes that cleave heme to produce the antioxidant biliverdin, inorganic iron and carbon monoxide (CO). The activity of Heme Oxygenase 1 (HO-1), also designated HSP 32, is highly inducible in response to numerous stimuli, including heme, heavy metals, hormones and oxidative stress. Heme Oxygenase 2, in contrast, appears to be constitutively expressed in mammalian tissues. Heme Oxygenase 2 is involved in the production of carbon monoxide (CO) in brain, where CO is thought to act as a neurotransmitter. The CO signaling system closely parallels the signaling pathway involving nitric oxide, and regulation of the two systems is closely linked. Heme Oxygenase 3 is found in the spleen, liver, thymus, prostate, heart, kidney, brain and testis. A poor heme catalyst, Heme Oxygenase 3 has two heme regulatory motifs that may be involved in heme binding.

REFERENCES

1. Maines, M.D. 1988. Heme oxygenase: function, multiplicity, regulatory mechanisms, and clinical applications. *FASEB J.* 2: 2557-2568.
2. Rodgers, P.A., et al. 1990. Developmental biology of heme oxygenase. *Clin. Perinatol.* 17: 275-291.
3. Alam, J., et al. 1994. Isolation and characterization of the mouse heme oxygenase 1 gene. Distal 5' sequences are required for induction by heme or heavy metals. *J. Biol. Chem.* 269: 1001-1009.
4. Maines, M.D. 1997. The heme oxygenase system; a regulator of second messenger gases. *Annu. Rev. Pharmacol. Toxicol.* 37: 517-554.

CHROMOSOMAL LOCATION

Genetic locus: HMOX1 (human) mapping to 22q12.3.

PRODUCT

Heme Oxygenase 1 siRNA (h2) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Heme Oxygenase 1 shRNA Plasmid (h2): sc-44306-SH and Heme Oxygenase 1 shRNA (h2) Lentiviral Particles: sc-44306-V as alternate gene silencing products.

For independent verification of Heme Oxygenase 1 (h2) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44306A, sc-44306B and sc-44306C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Heme Oxygenase 1 siRNA (h2) is recommended for the inhibition of Heme Oxygenase 1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Heme Oxygenase 1 (A-3): sc-136960 is recommended as a control antibody for monitoring of Heme Oxygenase 1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Heme Oxygenase 1 gene expression knockdown using RT-PCR Primer: Heme Oxygenase 1 (h2)-PR: sc-44306-PR (20 μ l, 564 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Choi, Y.K., et al. 2010. Carbon monoxide promotes VEGF expression by increasing HIF-1 α protein level via two distinct mechanisms, translational activation and stabilization of HIF-1 α protein. *J. Biol. Chem.* 285: 32116-32125.
2. Ohgari, Y., et al. 2011. Roles of porphyrin and iron metabolisms in the δ -aminolevulinic acid (ALA)-induced accumulation of protoporphyrin and photodamage of tumor cells. *Photochem. Photobiol.* 87: 1138-1145.
3. Lin, P.H., et al. 2013. TRC8 suppresses tumorigenesis through targeting Heme Oxygenase 1 for ubiquitination and degradation. *Oncogene* 32: 2325-2334.

RESEARCH USE

For research use only, not for use in diagnostic procedures.