

RBP siRNA (m): sc-44578

BACKGROUND

Retinol (Vitamin A) is transported in the blood bound to its carrier protein, retinol-binding protein (RBP), also designated plasma retinol-binding protein (PRBP) or RBP4. A member of the lipocalin family, RBP conveys retinol from stores in the liver to peripheral tissues. In plasma, RBP binds transthyretin (TTR, formerly called prealbumin) to prevent glomerular filtration of low molecular weight RBP in the kidneys. The stability of this complex holds diagnostic importance because the molar ratio of RBP:TTR provides an indirect way to indicate marginal vitamin A deficiency. Vitamin A deficiency blocks the secretion of RBP resulting in defective delivery and supply to epidermal cells. Originally identified solely as a transporter protein, recent studies correlating increased levels of RBP expression in adipose tissue with Insulin resistance have generated research into the possible roles the protein may play in the pathogenesis of type 2 diabetes and obesity.

REFERENCES

1. van Bennekum, A.M., et al. 1993. Retinol uptake from retinol-binding protein (RBP) by liver parenchymal cells *in vitro* does not specifically depend on its binding to RBP. *Biochemistry* 32: 1727-1733.
2. Zanotti, G., et al. 1993. The interaction of N-ethyl retinamide with plasma retinol-binding protein (RBP) and the crystal structure of the retinoid-RBP complex at 1.9-Å resolution. *J. Biol. Chem.* 268: 24873-24879.
3. Yamamoto, Y., et al. 1997. Interactions of transthyretin (TTR) and retinol-binding protein (RBP) in the uptake of retinol by primary rat hepatocytes. *Exp. Cell Res.* 234: 373-378.
4. Naylor, H.M., et al. 1999. The structure of human retinol-binding protein (RBP) with its carrier protein transthyretin reveals an interaction with the carboxy terminus of RBP. *Biochemistry* 38: 2647-2653.
5. Quadro, L., et al. 2002. Muscle expression of human retinol-binding protein (RBP). Suppression of the visual defect of RBP knockout mice. *J. Biol. Chem.* 277: 30191-30197.
6. Monaco H.L. 2002. Three-dimensional structure of the transthyretin-retinol-binding protein complex. *Clin. Chem. Lab. Med.* 40: 1229-1236.
7. Rosales, F.J., et al. 2002. Determination of a cut-off value for the molar ratio of retinol-binding protein to transthyretin (RBP:TTR) in Bangladeshi patients with low hepatic vitamin A stores. *J. Nutr.* 132: 3687-3692.
8. Raghu, P., et al. 2003. A new method for purification of human plasma retinol-binding protein and transthyretin. *Biotechnol. Appl. Biochem.* 38: 19-24.
9. Yang, Q., et al. 2005. Serum retinol binding protein 4 contributes to insulin resistance in obesity and type 2 diabetes. *Nature* 436: 356-362.

CHROMOSOMAL LOCATION

Genetic locus: Rbp4 (mouse) mapping to 19 C3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

RBP siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RBP shRNA Plasmid (m): sc-44578-SH and RBP shRNA (m) Lentiviral Particles: sc-44578-V as alternate gene silencing products.

For independent verification of RBP (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44578A, sc-44578B and sc-44578C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RBP siRNA (m) is recommended for the inhibition of RBP expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

RBP (20F9): sc-69795 is recommended as a control antibody for monitoring of RBP gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RBP gene expression knockdown using RT-PCR Primer: RBP (m)-PR: sc-44578-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.