

Dynein IC2, cytosolic siRNA (h): sc-44683

BACKGROUND

Dyneins are multisubunit, high molecular weight ATPases that interact with microtubules to generate force by converting the chemical energy of ATP into the mechanical energy of movement. Cytoplasmic or axonemal Dynein heavy, intermediate, light and light-intermediate chains are all components of minus end-directed motors; the complex transports cellular cargos towards the central region of the cell. Axonemal Dynein motors contain one to three non-identical heavy chains and cause a sliding of microtubules in the axonemes of cilia and flagella in a mechanism necessary for cilia to beat and propel the cell. Cytoplasmic Dynein is an approximately 12 subunit complex of 2 heavy chains, 2 intermediate chains to anchor Dynein to its cargo, four smaller intermediate chains and several light chains. It performs functions necessary for cell survival such as organelle transport and centrosome assembly. The carboxy terminus of Dynein is important for microtubule-dependent motility and is highly conserved, while the amino terminal regions are more variable. Several proteins regulate Dynein activity, including dynactin, LIS1 and NudEL(NudE-like).

REFERENCES

1. Mallik, R., et al. 2004. Cytoplasmic Dynein functions as a gear in response to load. *Nature* 427: 649-652.
2. Malikov, V., et al. 2004. Cytoplasmic Dynein nucleates microtubules to organize them into radial arrays *in vivo*. *Mol. Biol. Cell* 15: 2742-2749.
3. Asai, D.J., et al. 2004. The Dynein heavy chain family. *J. Eukaryot. Microbiol.* 51: 23-29.
4. Li, J., et al. 2005. NudEL targets Dynein to microtubule ends through LIS1. *Nat. Cell Biol.* 7: 686-690.
5. Seetharam, R.N., et al. 2005. High speed sliding of axonemal microtubules produced by outer arm Dynein. *Cell Motil. Cytoskeleton* 60: 96-103.
6. McGrath, J.L. 2005. Dynein motility: four heads are better than two. *Curr. Biol.* 15: R970-R972.
7. Lee, W.L., et al. 2005. The offloading model for Dynein function: differential function of motor subunits. *J. Cell Biol.* 168: 201-207.
8. He, Y., et al. 2005. Role of cytoplasmic Dynein in the axonal transport of microtubules and neurofilaments. *J. Cell Biol.* 168: 697-703.
9. Pfister, K.K., et al. 2005. Cytoplasmic Dynein nomenclature. *J. Cell Biol.* 171: 411-413.

CHROMOSOMAL LOCATION

Genetic locus: DYNC1H2 (human) mapping to 2q31.1.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Dynein IC2, cytosolic siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Dynein IC2, cytosolic shRNA Plasmid (h): sc-44683-SH and Dynein IC2, cytosolic shRNA (h) Lentiviral Particles: sc-44683-V as alternate gene silencing products.

For independent verification of Dynein IC2, cytosolic (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44683A, sc-44683B and sc-44683C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Dynein IC2, cytosolic siRNA (h) is recommended for the inhibition of Dynein IC2, cytosolic expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Dynein IC2, cytosolic gene expression knockdown using RT-PCR Primer: Dynein IC2, cytosolic (h)-PR: sc-44683-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.