

GGTase-II β siRNA (h): sc-45437

BACKGROUND

Eukaryotic cells contain three distinct prenyltransferases that catalyze the attachment of a thioether-linked 15-carbon farnesyl group or 20-carbon geranylgeranyl group to C-terminal cysteine residues. Geranylgeranyltransferase type I (GGTase-I, PGGTase-I) catalyzes the nucleophilic substitution reaction between geranylgeranyl diphosphate (GGPP) and a protein-derived thiol to form the thioether linkage. The candidate protein contains a C-terminal CAAX motif in which "A" is an aliphatic amino acid and "X" is leucine. Geranylgeranylation is necessary for the TGF β 1 signaling pathway, which involves phosphatidylcholine-specific phospholipase and a protein kinase C. Human GGTase-I contains an α subunit and a β subunit. Geranylgeranyltransferase type II (GGTase-II) is a heterodimer that catalyzes the transfer of two 20-carbon geranylgeranyl groups from geranylgeranyl pyrophosphate onto C-terminal cysteine residues of Rab GTPases, which is required for the activity of Rab proteins. GGTase-II also contains an α subunit and a β subunit.

REFERENCES

1. Schafer, W.R., et al. 1992. Protein prenylation: genes, enzymes, targets, and functions. *Annu. Rev. Genet.* 26: 209-237.
2. van Bokhoven, H., et al. 1996. cDNA cloning and chromosomal localization of the genes encoding the α - and β -subunits of human Rab geranylgeranyl transferase: the 3' end of the α -subunit gene overlaps with the transglutaminase 1 gene promoter. *Genomics* 38: 133-140.
3. Online Mendelian Inheritance in Man, OMIM[™]. 1997. Johns Hopkins University, Baltimore, MD. MIM Number: 602031. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Desnoyers, L., et al. 1998. Single prenyl-binding site on protein prenyl transferases. *Proc. Natl. Acad. Sci. USA* 95: 12266-12270.
5. Song, H.J., et al. 1998. Requirement for geranylgeranyl transferase I and acyl transferase in the TGF- β -stimulated pathway leading to elastin mRNA stabilization. *Biochem. Biophys. Res. Commun.* 252: 111-116.
6. Kalinin, A., et al. 2001. Expression of mammalian geranylgeranyltransferase type-II in *Escherichia coli* and its application for *in vitro* prenylation of Rab proteins. *Protein Expr. Purif.* 22: 84-91.

CHROMOSOMAL LOCATION

Genetic locus: RABGGTB (human) mapping to 1p31.1.

PRODUCT

GGTase-II β siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GGTase-II β shRNA Plasmid (h): sc-45437-SH and GGTase-II β shRNA (h) Lentiviral Particles: sc-45437-V as alternate gene silencing products.

For independent verification of GGTase-II β (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45437A, sc-45437B and sc-45437C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GGTase-II β siRNA (h) is recommended for the inhibition of GGTase-II β expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

GGTase-II β (B-8): sc-365926 is recommended as a control antibody for monitoring of GGTase-II β gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GGTase-II β gene expression knockdown using RT-PCR Primer: GGTase-II β (h)-PR: sc-45437-PR (20 μ l, 535 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.