



NMNAT-1 siRNA (m): sc-45503

BACKGROUND

Nicotinamide adenine dinucleotide (NMNAT) is an essential cofactor involved in fundamental processes in cell metabolism. NMNAT plays a key role in NAD⁺ biosynthesis, catalyzing the condensation of nicotinamide mononucleotide and ATP and yielding NAD⁺ and pyrophosphate. NMNAT appears to be a substrate of nuclear kinases and contains at least three potential phosphorylation sites. The interaction of NMNAT with nuclear proteins is likely to be modulated by phosphorylation. NMNAT is widely expressed with highest levels in skeletal muscle, heart, liver and kidney.

REFERENCES

1. D'Angelo, I., et al. 2000. Structure of nicotinamide mononucleotide adenyltransferase: a key enzyme in NAD⁺ biosynthesis. *Structure* 8: 993-1004.
2. Schweiger, M., et al. 2001. Characterization of recombinant human nicotinamide mononucleotide adenyltransferase (NMNAT), a nuclear enzyme essential for NAD synthesis. *FEBS Lett.* 492: 95-100.
3. Mack, T.G., et al. 2001. Wallerian degeneration of injured axons and synapses is delayed by a Ube4b/NMNAT chimeric gene. *Nat. Neurosci.* 4: 1199-1206.
4. Werner, E., et al. 2002. Crystallization and preliminary X-ray analysis of human nicotinamide mononucleotide adenyltransferase (NMNAT). *Acta Crystallogr. D Biol. Crystallogr.* 58: 140-142.
5. Gillingwater, T.H., et al. 2002. Age-dependent synapse withdrawal at axotomized neuromuscular junctions in Wld(s) mutant and Ube4b/NMNAT transgenic mice. *J. Physiol.* 543: 739-755.

CHROMOSOMAL LOCATION

Genetic locus: Nmnat1 (mouse) mapping to 4 E2.

PRODUCT

NMNAT-1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NMNAT-1 shRNA Plasmid (m): sc-45503-SH and NMNAT-1 shRNA (m) Lentiviral Particles: sc-45503-V as alternate gene silencing products.

For independent verification of NMNAT-1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45503A, sc-45503B and sc-45503C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NMNAT-1 siRNA (m) is recommended for the inhibition of NMNAT-1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NMNAT-1 gene expression knockdown using RT-PCR Primer: NMNAT-1 (m)-PR: sc-45503-PR (20 μ l, 457 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Nam, T.S., et al. 2020. Interleukin-8 drives CD38 to form NAADP from NADP⁺ and NAAD in the endolysosomes to mobilize Ca²⁺ and effect cell migration. *FASEB J.* 34: 12565-12576.
2. Dong, H., et al. 2024. Nuclear nicotinamide adenine dinucleotide deficiency by Nmnat1 deletion impaired hepatic insulin signaling, mitochondrial function, and hepatokine expression in mice fed a high-fat diet. *Lab. Invest.* 104: 100329.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.