

Lipophilin B siRNA (h): sc-45744

BACKGROUND

Secretoglobins (also called lipophilins or mammaglobins) are small secreted proteins of endocrine-responsive organs and mucosal epithelia that form multimeric complexes and correlate with the development of various human cancers. Lipophilin A and Lipophilin B are orthologs of prostatein (estradiol-binding protein), the major secretory glycoprotein of the rat ventral prostate gland. Lipophilin A, also designated LIPA, LPHA and secretoglobulin, family 1D, member 1 (SCGB1D1), is a component of a heterodimeric molecule present in human tears. Lipophilin B, also designated LIPB, LPHB and secretoglobulin, family 1D, member 2 (SCGB1D2), mRNA can be overexpressed in breast tumors and shows a high degree of correlation with the mRNA expression profile of mammaglobin. Histological detection in breast tissue of Mammaglobin A, also designated MGB1 and secretoglobulin, family 2A, member 2 (SCGB2A2) and Mammaglobin B, also designated MGB2, Lipophilin C, LPHC, UGB3 and SCGB2A2, is a reliable diagnostic marker for breast tumors.

REFERENCES

1. Wood, D.D., et al. 1984. Interaction between human Myelin basic protein and Lipophilin. *Neurochem. Res.* 9: 1523-1531.
2. Gow, A. 1997. Redefining the Lipophilin family of proteolipid proteins. *J. Neurosci. Res.* 50: 659-664.
3. Lehrer, R.I., et al. 1998. Lipophilin, a novel heterodimeric protein of human tears. *FEBS Lett.* 432: 163-167.
4. Carter, D., et al. 2002. Purification and characterization of the mammaglobin/lipophilin B complex, a promising diagnostic marker for breast cancer. *Biochemistry* 41: 6714-6722.
5. Carter, D., et al. 2003. Serum antibodies to Lipophilin B detected in late stage breast cancer patients. *Clin. Cancer Res.* 9: 749-754.
6. Cerveira, N., et al. 2004. Highly sensitive detection of the MGB1 transcript (Mammaglobin) in the peripheral blood of breast cancer patients. *Int. J. Cancer* 108: 592-595.
7. Ouellette, R.J., et al. 2004. RT-PCR for Mammaglobin genes, MGB1 and MGB2, identifies breast cancer micrometastases in sentinel lymph nodes. *Am. J. Clin. Pathol.* 121: 637-643.
8. Sjodin, A., et al. 2005. Secretoglobins in the human pituitary: high expression of lipophilin B and its down-regulation in pituitary adenomas. *Acta Neuropathol.* 109: 381-386.
9. Bernstein, J.L., et al. 2005. Identification of mammaglobin as a novel serum marker for breast cancer. *Clin. Cancer Res.* 11: 6528-6535.

CHROMOSOMAL LOCATION

Genetic locus: SCGB1D1 (human) mapping to 11q12.3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Lipophilin A siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Lipophilin A shRNA Plasmid (h): sc-45743-SH and Lipophilin A shRNA (h) Lentiviral Particles: sc-45743-V as alternate gene silencing products.

For independent verification of Lipophilin A (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45743A, sc-45743B and sc-45743C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Lipophilin A siRNA (h) is recommended for the inhibition of Lipophilin A expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RESEARCH USE

For research use only, not for use in diagnostic procedures.