

# $G_{\alpha 11}$ siRNA (r): sc-45999

## BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (a photon, pheromone, odorant, hormone or neurotransmitter) while the effectors (e.g., adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. In mammals, G protein  $\alpha$ ,  $\beta$  and  $\gamma$  polypeptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their  $\alpha$  subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity of the best studied effectors. Four distinct classes of  $G_{\alpha}$  subunits have been identified; these include  $G_s$ ,  $G_i$ ,  $G_q$  and  $G_{12/13}$ . The  $G_q$  class includes  $G_{\alpha 15}$ ,  $G_{\alpha 14}$ ,  $G_{\alpha 11}$  and  $G_{\alpha q}$ , two of which,  $G_{\alpha 11}$  and  $G_{\alpha q}$ , are abundant in brain and lung and present at lower levels in a variety of tissues.

## REFERENCES

1. Strathmann, M. and Simon, M.I. 1990. G Protein diversity: a distinct class of  $\alpha$  subunits is present in vertebrates and invertebrates. *Proc. Natl. Acad. Sci. USA* 87: 9113-9117.
2. Simon, M.I., Strathmann, M.P. and Gautam, N. 1991. Diversity of G proteins in signal transduction. *Science* 252: 802-808.
3. Cali, J.J., Balcueva, E.A., Rybalkin, I. and Robishaw, J.D. 1992. Selective tissue distribution of G protein  $\gamma$  subunits, including a new form of the  $\gamma$  subunits identified by cDNA cloning. *J. Biol. Chem.* 267: 24023-24027.
4. McLaughlin, S.K., McKinnon, P.J. and Margolskee, R.F. 1992. Gustducin is a taste-cell-specific G protein closely related to the transducins. *Nature* 357: 563-569.
5. von Weizsäcker, E., Strathman, M.P. and Simon, M.I. 1992. Diversity among the  $\beta$  subunits of heterotrimeric GTP-binding proteins: characterization of a novel  $\beta$  subunit cDNA. *Biochem. Biophys. Res. Commun.* 183: 350-356.
6. Conklin, B.R. and Bourne, H.R. 1993. Structural elements of  $G_{\alpha}$  subunits that interact with  $G_{\beta\gamma}$  receptors and effectors. *Cell* 73: 631-641.

## CHROMOSOMAL LOCATION

Genetic locus: Gna11 (rat) mapping to 7q11.

## PRODUCT

$G_{\alpha 11}$  siRNA (r) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see  $G_{\alpha 11}$  shRNA Plasmid (r): sc-45999-SH and  $G_{\alpha 11}$  shRNA (r) Lentiviral Particles: sc-45999-V as alternate gene silencing products.

For independent verification of  $G_{\alpha 11}$  (r) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-45999A, sc-45999B and sc-45999C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

$G_{\alpha 11}$  shRNA Plasmid (r) is recommended for the inhibition of  $G_{\alpha 11}$  expression in rat cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

$G_{\alpha 11}$  (D-6): sc-390382 is recommended as a control antibody for monitoring of  $G_{\alpha 11}$  gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor  $G_{\alpha 11}$  gene expression knockdown using RT-PCR Primer:  $G_{\alpha 11}$  (r)-PR: sc-45999-PR (20  $\mu$ l, 580 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.