

survivin (8E2): sc-47750

BACKGROUND

The baculovirus protein p35 inhibits virally-induced apoptosis of invertebrate and mammalian cells and may function to impair the clearing of virally infected cells by the immune system of the host. This is accomplished at least in part by the ability of p35 to block both TNF- and FAS-mediated apoptosis through the inhibition of the ICE family of serine proteases. Two mammalian homologs of baculovirus p35, referred to as inhibitor of apoptosis protein (IAP) 1 and 2, share an amino-terminal baculovirus IAP repeat (BIR) motif and a carboxy-terminal RING finger. Although the c-IAPs do not directly associate with the TNF receptor (TNF-R), they efficiently block TNF-mediated apoptosis through their interaction with the downstream TNF-R effectors, TRAF1 and TRAF2. Additional IAP family members include ILP (for IAP-like protein) and survivin. ILP inhibits activated caspase-3, leading to the resistance of FAS-mediated apoptosis. Survivin (also designated TIAP) is expressed during the G₂/M phase of the cell cycle and associates with microtubules of the mitotic spindle. Increased caspase-3 activity is detected when a disruption of survivin-microtubule interactions occurs.

CHROMOSOMAL LOCATION

Genetic locus: BIRC5 (human) mapping to 17q25.3; Birc5 (mouse) mapping to 11 E2.

SOURCE

survivin (8E2) is a mouse monoclonal antibody raised against full length human survivin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

survivin (8E2) is recommended for detection of survivin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for survivin siRNA (h): sc-29499, survivin siRNA (m): sc-29500, survivin shRNA Plasmid (h): sc-29499-SH, survivin shRNA Plasmid (m): sc-29500-SH, survivin shRNA (h) Lentiviral Particles: sc-29499-V and survivin shRNA (m) Lentiviral Particles: sc-29500-V.

Molecular Weight of survivin: 17 kDa.

Positive Controls: mouse spleen extract: sc-2391, rat heart extract: sc-2393 or rat lung extract: sc-2396.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



survivin (8E2): sc-47750. Western blot analysis of survivin expression in mouse spleen (A) and rat heart (B) tissue extracts.



survivin (8E2): sc-47750. Western blot analysis of survivin expression in rat lung tissue extract.

SELECT PRODUCT CITATIONS

- Gao, Y., et al. 2010. N-terminal deletion effects of human survivin on dimerization and binding to Smac/DIABLO *in vitro*. J. Phys. Chem. B 114: 15656-15662.
- Yu, R.X., et al. 2011. Effects of fucoxanthin on proliferation and apoptosis in human gastric adenocarcinoma MGC-803 cells via JAK/STAT signal pathway. Eur. J. Pharmacol. 657: 10-19.
- Rajendran, P., et al. 2011. Suppression of signal transducer and activator of transcription 3 activation by butein inhibits growth of human hepatocellular carcinoma *in vivo*. Clin. Cancer Res. 17: 1425-1439.
- Wan, X.B., et al. 2012. Molecular prognostic prediction for locally advanced nasopharyngeal carcinoma by support vector machine integrated approach. PLoS ONE 7: e31989.
- Wang, J., et al. 2012. *In vivo* induction of apoptosis by fucoxanthin, a marine carotenoid, associated with down-regulating STAT3/EGFR signaling in sarcoma 180 (S180) xenografts-bearing mice. Mar. Drugs 10: 2055-2068.
- Malhotra, U., et al. 2013. Prognostic value and targeted inhibition of survivin expression in esophageal adenocarcinoma and cancer-adjacent squamous epithelium. PLoS ONE 8: e78343.
- Zheng, W.X., et al. 2014. Baicalin protects PC-12 cells from oxidative stress induced by hydrogen peroxide via anti-apoptotic effects. Brain Inj. 28: 227-234.
- Cai, H., et al. 2019. Molecular decision tree algorithms predict individual recurrence pattern for locally advanced nasopharyngeal carcinoma. J. Cancer 10: 3323-3332.



See **survivin (D-8): sc-17779** for survivin antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.