

Integrin $\alpha 2$ (244G1): sc-47770

BACKGROUND

Integrins are heterodimers composed of noncovalently associated transmembrane α and β subunits. The 16 α and 8 β subunits heterodimerize to produce more than 20 different receptors. Most integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, collagen and Vitronectin. Certain integrins can also bind to soluble ligands such as fibrinogen, or to counter-receptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster integrins by binding to adjacent integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis. Integrin $\alpha 2$ is responsible for adhesion of platelets and other cells to collagens. Modulation of collagen and collagenase gene expression force generation and organization of newly synthesized extracellular matrix.

REFERENCES

1. Takada, Y. and Hemler, M. 1989. The primary structure of the VLA-2/collagen receptor $\alpha 2$ subunit (platelet GPIa): homology to other integrins and the presence of a possible collagen-binding domain. *J. Cell Biol.* 109: 397-407.
2. Hynes, R.O. 1992. Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 69: 11-25.
3. Santoso, S., et al. 1993. The human platelet alloantigens Br(a) and Br(b) are associated with a single amino acid polymorphism on glycoprotein Ia (integrin subunit $\alpha 2$). *J. Clin. Invest.* 92: 2427-2432.
4. Miyamoto, S., et al. 1995. Synergistic roles for receptor occupancy and aggregation in integrin transmembrane function. *Science* 267: 883-885.
5. Sheppard, D. 1996. Epithelial integrins. *Bioessays* 18: 655-660.
6. Juliano, R. 1996. Cooperation between soluble factors and integrin-mediated cell anchorage in the control of cell growth and differentiation. *Bioessays* 18: 911-917.
7. Moore, T.A., et al. 1996. Inhibition of $\gamma \delta$ T cell development and early thymocyte maturation in IL-7^{-/-} mice. *J. Immunol.* 157: 2366-2373.
8. Kroll, H., et al. 2000. The impact of the glycoprotein Ia collagen receptor subunit A1648G gene polymorphism on coronary artery disease and acute myocardial infarction. *Thromb. Haemost.* 83: 392-396.
9. Arase, H., et al. 2001. Cutting edge: the mouse NK cell-associated antigen recognized by DX5 monoclonal antibody is CD49b ($\alpha 2$ integrin, very late antigen-2). *J. Immunol.* 167: 1141-1144.

CHROMOSOMAL LOCATION

Genetic locus: ITGA2 (human) mapping to 5q11.2.

SOURCE

Integrin $\alpha 2$ (244G1) is a mouse monoclonal antibody raised against recombinant Integrin $\alpha 2$ of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS


Integrin $\alpha 2$ (244G1) is recommended for detection of Integrin $\alpha 2$ of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1×10^6 cells).

Suitable for use as control antibody for Integrin $\alpha 2$ siRNA (h): sc-29371, Integrin $\alpha 2$ shRNA Plasmid (h): sc-29371-SH and Integrin $\alpha 2$ shRNA (h) Lentiviral Particles: sc-29371-V.


Molecular Weight of Integrin $\alpha 2$: 150 kDa.

Positive Controls: HCT-116 whole cell lysate: sc-364175, MDA-MB-231 cell lysate: sc-2232 or WiDr cell lysate: sc-24779.

DATA



Integrin $\alpha 2$ (244G1): sc-47770. Western blot analysis of Integrin $\alpha 2$ expression in MDA-MB-231 (A) and WiDr (B) whole cell lysates.



Integrin $\alpha 2$ (244G1): sc-47770. Western blot analysis of Integrin $\alpha 2$ expression in HCT-116 whole cell lysate under non-reducing conditions.

SELECT PRODUCT CITATIONS

1. Wang, C., et al. 2020. Secreted pyruvate kinase M2 promotes lung cancer metastasis through activating the Integrin $\beta 1$ /FAK signaling pathway. *Cell Rep.* 30: 1780-1797.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.