

HDAC4 (D-1): sc-48390

BACKGROUND

In the intact cell, DNA closely associates with histones and other nuclear proteins to form chromatin. The remodeling of chromatin is believed to be a critical component of transcriptional regulation, and a major source of this remodeling is brought about by the acetylation of nucleosomal histones. Acetylation of lysine residues in the amino-terminal tail domain of histone results in an allosteric change in the nucleosomal conformation and an increased accessibility to transcription factors by DNA. Conversely, the deacetylation of histones is associated with transcriptional silencing. Several mammalian proteins have been identified as nuclear histone acetylases, including GCN5, p300/CBP, PCAF (p300/CBP-associated factor), HAT1 and the TFIID subunit TAF II p250. Mammalian HDAC1 (also designated HD1), HDAC2 (also designated RPD3) and HDAC3-6 have been identified as histone deacetylases.

REFERENCES

1. Lee, D.Y., et al. 1993. A positive role for histone acetylation in transcription factor access to nucleosomal DNA. *Cell* 72: 73-82.
2. Braunstein, M., et al. 1993. Transcriptional silencing in yeast is associated with reduced nucleosome acetylation. *Genes Dev.* 7: 592-604.

CHROMOSOMAL LOCATION

Genetic locus: HDAC4 (human) mapping to 2q37.3; Hdac4 (mouse) mapping to 1 D.

SOURCE

HDAC4 (D-1) is a mouse monoclonal antibody raised against amino acids 530-631 of HDAC4 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

HDAC4 (D-1) is recommended for detection of HDAC4 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for HDAC4 siRNA (h): sc-35540, HDAC4 siRNA (m): sc-35541, HDAC4 shRNA Plasmid (h): sc-35540-SH, HDAC4 shRNA Plasmid (m): sc-35541-SH, HDAC4 shRNA (h) Lentiviral Particles: sc-35540-V and HDAC4 shRNA (m) Lentiviral Particles: sc-35541-V.


Molecular Weight of HDAC4: 140 kDa.

Positive Controls: COLO 205 whole cell lysate: sc-364177, SUP-T1 whole cell lysate: sc-364796 or TK-1 whole cell lysate: sc-364798.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



HDAC4 (D-1): sc-48390. Western blot analysis of HDAC4 expression in COLO 205 (A), CCRF-CEM (B), SUP-T1 (C), TK-1 (D) and Neuro-2A (E) whole cell lysates.



HDAC4 (D-1): sc-48390. Immunoperoxidase staining of formalin fixed, paraffin-embedded human breast tissue showing cytoplasmic staining of glandular cells.

SELECT PRODUCT CITATIONS

1. Bhandari, D.R., et al. 2011. The regulatory role of c-MYC on HDAC2 and PcG expression in human multipotent stem cells. *J. Cell. Mol. Med.* 15: 1603-1614.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **HDAC4 (A-4): sc-46672** for HDAC4 antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647.