

NURIM (E-11): sc-514010



The Power to Question

BACKGROUND

NURIM, also known as NRM or NRM29, is a 262 amino acid multi-pass membrane protein that localizes to the inner membrane of the nucleus. Existing as multiple alternatively spliced isoforms, NURIM is thought to possess enzymatic functions that may play a role in nuclear envelope (NE) dynamics. The gene encoding NURIM maps to human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Additionally, porphyria cutanea tarda, Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder are all associated with genes that map to chromosome 6.

REFERENCES

1. Rolls, M.M., et al. 1999. A visual screen of a GFP-fusion library identifies a new type of nuclear envelope membrane protein. *J. Cell Biol.* 146: 29-44.
2. Holmer, L. and Worman, H.J. 2001. Inner nuclear membrane proteins: functions and targeting. *Cell. Mol. Life Sci.* 58: 1741-1747.
3. Otsuki, T., et al. 2005. Signal sequence and keyword trap in silico for selection of full-length human cDNAs encoding secretion or membrane proteins from oligo-capped cDNA libraries. *DNA Res.* 12: 117-126.
4. Hofemeister, H. and O'Hare, P. 2005. Analysis of the localization and topology of nurim, a polytopic protein tightly associated with the inner nuclear membrane. *J. Biol. Chem.* 280: 2512-2521.

CHROMOSOMAL LOCATION

Genetic locus: NRM (human) mapping to 6p21.33; Nrm (mouse) mapping to 17 B1.

SOURCE

NURIM (E-11) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 28-47 near the N-terminus of NURIM of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

NURIM (E-11) is available conjugated to agarose (sc-514010 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-514010 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-514010 PE), fluorescein (sc-514010 FITC), Alexa Fluor® 488 (sc-514010 AF488), Alexa Fluor® 546 (sc-514010 AF546), Alexa Fluor® 594 (sc-514010 AF594) or Alexa Fluor® 647 (sc-514010 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-514010 AF680) or Alexa Fluor® 790 (sc-514010 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-514010 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

NURIM (E-11) is recommended for detection of NURIM isoforms 1-5 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NURIM siRNA (h): sc-95631, NURIM siRNA (m): sc-150128, NURIM shRNA Plasmid (h): sc-95631-SH, NURIM shRNA Plasmid (m): sc-150128-SH, NURIM shRNA (h) Lentiviral Particles: sc-95631-V and NURIM shRNA (m) Lentiviral Particles: sc-150128-V.


Molecular Weight of NURIM: 30 kDa.

Positive Controls: NURIM (m2): 293T Lysate: sc-122182, PC-3 cell lysate: sc-2220 or IMR-32 nuclear extract: sc-2148.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



NURIM (E-11): sc-514010. Western blot analysis of NURIM expression in IMR-32 (A), SH-SY5Y (B), SK-N-SH (C), Neuro-2A (D) and NIH/3T3 (E) whole cell lysates and mouse brain tissue extract (F).



NURIM (E-11): sc-514010. Western blot analysis of NURIM expression in non-transfected 293T: sc-117752 (A), mouse NURIM transfected 293T: sc-122182 (B) and PC-3 (C) whole cell lysates, PC-3 (D) and IMR-32 (E) nuclear extracts and human testis tissue extract (F).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.