

## RBM12 (B-12): sc-514258



The Power to Question

## BACKGROUND

Proteins containing RNA recognition motifs, including various hnRNP proteins, are implicated in the regulation of alternative splicing and protein components of snRNPs. The RBM (RNA-binding motif) gene family encodes proteins with an RNA binding motif that have been suggested to play a role in the modulation of apoptosis. RBM12 (RNA binding motif protein 12), also known as SWAN, HRIHFB2091 or KIAA0765, is a 932 amino acid protein which localizes to the nucleus. RBM12 contains multiple proline-rich regions, transmembrane domains and three RNA recognition motifs (RRM). It has been found that the genes for RBM12 and copine I (CPNE1) overlap at human chromosome location 20q11.22, sharing the promoter region and a 5'UTR (which are conserved in human, zebrafish and mouse), suggesting that a functional interaction between the two genes may exist.

## CHROMOSOMAL LOCATION

Genetic locus: RBM12 (human) mapping to 20q11.22; Rbm12 (mouse) mapping to 2 H1.

## SOURCE

RBM12 (B-12) is a mouse monoclonal antibody raised against amino acids 422-530 mapping within an internal region of RBM12 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RBM12 (B-12) is available conjugated to agarose (sc-514258 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-514258 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-514258 PE), fluorescein (sc-514258 FITC), Alexa Fluor® 488 (sc-514258 AF488), Alexa Fluor® 546 (sc-514258 AF546), Alexa Fluor® 594 (sc-514258 AF594) or Alexa Fluor® 647 (sc-514258 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-514258 AF680) or Alexa Fluor® 790 (sc-514258 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

## APPLICATIONS

RBM12 (B-12) is recommended for detection of RBM12 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RBM12 siRNA (h): sc-76364, RBM12 siRNA (m): sc-152725, RBM12 shRNA Plasmid (h): sc-76364-SH, RBM12 shRNA Plasmid (m): sc-152725-SH, RBM12 shRNA (h) Lentiviral Particles: sc-76364-V and RBM12 shRNA (m) Lentiviral Particles: sc-152725-V.

Molecular Weight (predicted) of RBM12: 97 kDa.


Molecular Weight (observed) of RBM12: 85 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, THP-1 cell lysate: sc-2238 or K-562 whole cell lysate: sc-2203.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



RBM12 (B-12): sc-514258. Western blot analysis of RBM12 expression in HeLa (A), THP-1 (B) and K-562 (C) whole cell lysates.



RBM12 (B-12): sc-514258. Immunofluorescence staining of methanol-fixed HeLa (A) and SW480 (B) cells showing nuclear localization.

## SELECT PRODUCT CITATIONS

- Steinberg, S., et al. 2017. Truncating mutations in RBM12 are associated with psychosis. *Nat. Genet.* 49: 1251-1254.
- Gao, C., et al. 2021. Increased RBM12 expression predicts poor prognosis in hepatocellular carcinoma based on bioinformatics. *J. Gastrointest. Oncol.* 12: 1905-1926.
- Semesta, K.M., et al. 2023. The psychosis risk factor RBM12 encodes a novel repressor of GPCR/cAMP signal transduction. *J. Biol. Chem.* 299: 105133.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA