

GATA-3 (C-11): sc-514427

BACKGROUND

Members of the GATA family share a conserved zinc finger DNA-binding domain and are capable of binding the WGATAR consensus sequence. GATA-1 is erythroid-specific and is responsible for the regulated transcription of erythroid genes. It is an essential component in the generation of the erythroid lineage. GATA-2 is expressed in embryonic brain and liver, HeLa and endothelial cells, as well as in erythroid cells. Studies with a modified GATA consensus sequence, AGATCTTA, have shown that GATA-2 and GATA-3 recognize this mutated consensus while GATA-1 has poor recognition of this sequence. This indicates broader regulatory capabilities of GATA-2 and GATA-3 than GATA-1. GATA-3 is highly expressed in T lymphocytes. GATA-4, GATA-5 and GATA-6 comprise a subfamily of transcription factors. Both GATA-4 and GATA-6 are found in heart, pancreas and ovary; lung and liver tissues exhibit GATA-6, but not GATA-4 expression. GATA-5 expression has been observed in differentiated heart and gut tissues and is present throughout the course of development in the heart. Although expression patterns of the various GATA transcription factors may overlap, it is not yet apparent how the GATA factors are able to discriminate in binding their appropriate target sites.

CHROMOSOMAL LOCATION

Genetic locus: GATA3 (human) mapping to 10p14; Gata3 (mouse) mapping to 2 A1.

SOURCE

GATA-3 (C-11) is a mouse monoclonal antibody raised against amino acids 167-214 of GATA-3 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-514427 X, 200 µg/0.1 ml.

APPLICATIONS

GATA-3 (C-11) is recommended for detection of GATA-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GATA-3 siRNA (h): sc-29331, GATA-3 siRNA (m): sc-35453, GATA-3 siRNA (r): sc-61845, GATA-3 shRNA Plasmid (h): sc-29331-SH, GATA-3 shRNA Plasmid (m): sc-35453-SH, GATA-3 shRNA Plasmid (r): sc-61845-SH, GATA-3 shRNA (h) Lentiviral Particles: sc-29331-V, GATA-3 shRNA (m) Lentiviral Particles: sc-35453-V and GATA-3 shRNA (r) Lentiviral Particles: sc-61845-V.

GATA-3 (C-11) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of GATA-3: 50 kDa.

Positive Controls: GATA-3 (h): 293 Lysate: sc-110481.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



GATA-3 (C-11): sc-514427. Western blot analysis of GATA-3 expression in non-transfected 293: sc-110760 (A), human GATA-3 transfected 293: sc-110481 (B), MOLT-4 (C) and CCRF-HSB-2 (D) whole cell lysates.



GATA-3 (C-11): sc-514427. Immunoperoxidase staining of formalin fixed, paraffin-embedded human breast tissue showing nuclear staining of glandular cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human urinary bladder tissue showing nuclear staining of urothelial cells (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **GATA-3 (HG3-31): sc-268** for GATA-3 antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647.