

CYP27A1 (D-2): sc-514693

BACKGROUND

P450 enzymes constitute a family of monooxygenase enzymes that are involved in the metabolism of a wide array of endogenous and xenobiotic compounds. P450 enzymes can be classified, based on their sequence similarities, into distinct subfamilies, which include CYP1A and CYP2A. Other P450 family members include CYP19, also designated aromatase (P450_{arom}), which catalyzes the conversion of C19 steroids to estrogens in various tissues, including placenta, gonads, adipose tissue, skin and brain. CYP19 expression is controlled by hormonally regulated promoters in different tissues and increased aromatase activity is associated with familial gynecomastia. Also, a polymorphic allele of CYP19 (repeat (TTTA)₁₂) is present in a majority of breast cancer patients. P450 cholesterol 7 α -hydroxylase, CYP7A1, is the rate limiting enzyme of bile acid synthesis in the liver, and its expression is mediated by the bile acid receptor FXR. CYP27A1 catalyzes vitamin D₃ 25-hydroxylation and is localized to the mitochondria in kidney and liver.

REFERENCES

- Nelson, D.R., et al. 1996. P450 superfamily: update on new sequences, gene mapping, accession numbers and nomenclature. *Pharmacogenetics* 6: 1-42.
- Peterson, J.A., et al. 1997. P450_{BM-3}; a tale of two domains—or is it three? *Steroids* 62: 117-123.
- Bulun, S.E., et al. 1997. Endocrine disorders associated with inappropriately high aromatase expression. *J. Steroid Biochem. Mol. Biol.* 61: 133-139.
- Braunstein, G.D. 1999. Aromatase and gynecomastia. *Endocr. Relat. Cancer* 6: 315-324.
- Kristensen, V.N., et al. 2000. Genetic variants of CYP19 (aromatase) and breast cancer risk. *Oncogene* 19: 1329-1333.
- Repa, J.J., et al. 2000. Regulation of absorption and ABC1-mediated efflux of cholesterol by RXR heterodimers. *Science* 289: 1524-1529.
- Sawada, N., et al. 2000. Metabolism of vitamin D₃ by human CYP27A1. *Biochem. Biophys. Res. Commun.* 273: 977-984.

CHROMOSOMAL LOCATION

Genetic locus: CYP27A1 (human) mapping to 2q35.

SOURCE

CYP27A1 (D-2) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 454-476 near the C-terminus of CYP27A1 of human origin.

PRODUCT

Each vial contains 200 μ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-514693 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

CYP27A1 (D-2) is recommended for detection of CYP27A1 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CYP27A1 siRNA (h): sc-41500, CYP27A1 shRNA Plasmid (h): sc-41500-SH and CYP27A1 shRNA (h) Lentiviral Particles: sc-41500-V.


Molecular Weight of CYP27A1: 60 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203 or A549 cell lysate: sc-2413.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



CYP27A1 (D-2): sc-514693. Western blot analysis of CYP27A1 expression in K-562 whole cell lysate.



CYP27A1 (D-2): sc-514693. Western blot analysis of CYP27A1 expression in A549 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.