

TFEC (E-12): sc-515031

BACKGROUND

The DNA-binding factor TFE3 contains adjacent helix-loop-helix (HLH) and leucine zipper (LZ) domains flanked by an upstream basic region. These protein motifs are frequently observed in other transcription factors and are particularly common to members of the Myc family. TFE3 is ubiquitously expressed and can directly associate with DNA as either homodimers or heterodimers formed with two related proteins, TFEB or TFEC. TFE3 binds to and activates the microE3 motif of the immunoglobulin heavy-chain enhancer to induce B cell-specific gene transcription and DNA recombination. TFEB binds to the major late promoter of adenovirus and specifically associates with DNA as both a homodimer and a heterodimer with TFE3. TFEB is expressed at low levels in the embryo but at high levels in the trophoblast cells of the placenta, where it plays a critical role in regulating normal vascularization of the placenta. TFEC shares a bHLH/LZ structure with TFE3 and a closely related protein microphthalmia-associated transcription factor (MITF), which is critically involved in melanocyte differentiation. Unlike TFE3, the expression of TFEC is largely restricted to fibroblasts, myoblasts, chondrosarcoma cells and myeloma cells.

REFERENCES

1. Beckmann, H., et al. 1990. TFE3: a helix-loop-helix protein that activates transcription through the immunoglobulin enhancer μ E3 motif. *Genes Dev.* 4: 167-179.
2. Fisher, D.E., et al. 1991. TFEB has DNA-binding and oligomerization properties of a unique helix-loop-helix/leucine-zipper family. *Genes Dev.* 5: 2342-2352.
3. Kerkhoff, E., et al. 1991. Sequence-specific DNA binding by Myc proteins. *Proc. Natl. Acad. Sci. USA* 88: 4323-4327.
4. Artandi, S.E., et al. 1994. The basic helix-loop-helix-zipper domain of TFE3 mediates enhancer-promoter interaction. *Mol. Cell. Biol.* 14: 7704-7716.

CHROMOSOMAL LOCATION

Genetic locus: TFEC (human) mapping to 7q31.2.

SOURCE

TFEC (E-12) is a mouse monoclonal antibody raised against amino acids 1-68 mapping at the N-terminus of TFEC of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

TFEC (E-12) is available conjugated to agarose (sc-515031 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-515031 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-515031 PE), fluorescein (sc-515031 FITC), Alexa Fluor[®] 488 (sc-515031 AF488), Alexa Fluor[®] 546 (sc-515031 AF546), Alexa Fluor[®] 594 (sc-515031 AF594) or Alexa Fluor[®] 647 (sc-515031 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-515031 AF680) or Alexa Fluor[®] 790 (sc-515031 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

TFEC (E-12) is recommended for detection of TFEC of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TFEC siRNA (h): sc-38511, TFEC shRNA Plasmid (h): sc-38511-SH and TFEC shRNA (h) Lentiviral Particles: sc-38511-V.


Molecular Weight of TFEC: 39 kDa.

Positive Controls: TFEC (h): 293T Lysate: sc-114466.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



TFEC (E-12): sc-515031. Western blot analysis of TFEC expression in non-transfected: sc-117752 (A) and human TFEC transfected: sc-114466 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

1. Kar, S.K., et al. 2020. Effects of undigested protein-rich ingredients on polarized small intestinal organoid monolayers. *J. Anim. Sci. Biotechnol.* 11: 51.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.