

NSF (A-3): sc-515043

BACKGROUND

Syntaxins were originally thought to be docking proteins, but have more recently been categorized as anchoring proteins that anchor themselves to the cytoplasmic surfaces of cellular membranes. Syntaxins have been shown to bind to various proteins involved in exocytosis, including VAMPs (vesicle-associated membrane proteins), NSF (N-ethylmaleimide-sensitive factor), SNAP 25 (synaptosomal-associated protein of 25 kDa), SNAPs (soluble NSF attachment proteins) and synaptotagmin. VAMPs, also designated synaptobrevins, including VAMP-1 and VAMP-2, and synaptotagmin, a protein that may function as an inhibitor of exocytosis, are vesicular proteins. SNAPs, including α - and γ -SNAP, are cytoplasmic proteins that bind to a membrane receptor complex composed of VAMP, SNAP 25 and syntaxin. SNAPs mediate the membrane binding of NSF, which is essential for membrane fusion reactions.

REFERENCES

1. Elferink, L.A., et al. 1993. A role for synaptotagmin (p65) in regulated exocytosis. *Cell* 72: 153-159.
2. Bennett, M.K., et al. 1993. The syntaxin family of vesicular transport receptors. *Cell* 74: 863-873.
3. Yamaguchi, K. and Akagawa, K. 1994. Exocytosis relating proteins in the nervous system. *Neurosci. Res.* 20: 289-292.
4. Hayashi, T., et al. 1994. Synaptic vesicle membrane fusion complex: action of clostridial neurotoxins on assembly. *EMBO J.* 13: 5051-5061.

CHROMOSOMAL LOCATION

Genetic locus: NSF (human) mapping to 17q21.31; Nsf (mouse) mapping to 11 E1.

SOURCE

NSF (A-3) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 50-69 near the N-terminus of NSF of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

NSF (A-3) is available conjugated to agarose (sc-515043 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-515043 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-515043 PE), fluorescein (sc-515043 FITC), Alexa Fluor[®] 488 (sc-515043 AF488), Alexa Fluor[®] 546 (sc-515043 AF546), Alexa Fluor[®] 594 (sc-515043 AF594) or Alexa Fluor[®] 647 (sc-515043 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-515043 AF680) or Alexa Fluor[®] 790 (sc-515043 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-515043 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

NSF (A-3) is recommended for detection of NSF of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NSF siRNA (h): sc-36101, NSF siRNA (m): sc-36102, NSF siRNA (r): sc-156016, NSF shRNA Plasmid (h): sc-36101-SH, NSF shRNA Plasmid (m): sc-36102-SH, NSF shRNA Plasmid (r): sc-156016-SH, NSF shRNA (h) Lentiviral Particles: sc-36101-V, NSF shRNA (m) Lentiviral Particles: sc-36102-V and NSF shRNA (r) Lentiviral Particles: sc-156016-V.


Molecular Weight of NSF: 76 kDa.

Positive Controls: SK-N-SH cell lysate: sc-2410, SK-N-MC cell lysate: sc-2237 or MIA PaCa-2 cell lysate: sc-2285.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



NSF (A-3): sc-515043. Western blot analysis of NSF expression in MIA PaCa-2 (A), BE (2)-M17 (B), H4 (C), Neuro-2A (D), C6 (E) and RiN-m5F (F) whole cell lysates.



NSF (A-3): sc-515043. Western blot analysis of NSF expression in SK-N-SH (A), SK-N-MC (B), MIA PaCa-2 (C), HeLa (D) and IMR-32 (E) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.