

N-Myc (C-4): sc-515099

BACKGROUND

The v-Myc oncogene, initially identified in the MC29 avian retrovirus, causes myelocytomas, carcinomas, sarcomas and lymphomas, and belongs to a family of oncogenes conserved throughout evolution. In humans, the family consists of five genes: c-Myc, N-Myc, R-Myc, L-Myc and B-Myc. Amplification of the N-Myc gene has been found in human neuroblastomas and cell lines. The extent of N-Myc amplification correlates well with the stage of neuroblastoma disease. Immunological studies have shown that the human N-Myc gene encodes a nuclear phosphoprotein that exhibits relatively short (30 min) half life *in vivo*. The prototype member of the family, c-Myc p67, binds DNA in a sequence-specific manner subsequent to dimerization with a second basic region helix-loop-helix leucine zipper motif protein, designated Max.

CHROMOSOMAL LOCATION

Genetic locus: MYCN (human) mapping to 2p24.3; Mycn (mouse) mapping to 12 A1.1.

SOURCE

N-Myc (C-4) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 443-464 within the C-terminus of N-Myc of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-515099 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-515099 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

N-Myc (C-4) is recommended for detection of N-Myc of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for N-Myc siRNA (h): sc-36003, N-Myc siRNA (m): sc-38087, N-Myc shRNA Plasmid (h): sc-36003-SH, N-Myc shRNA Plasmid (m): sc-38087-SH, N-Myc shRNA (h) Lentiviral Particles: sc-36003-V and N-Myc shRNA (m) Lentiviral Particles: sc-38087-V.

N-Myc (C-4) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of N-Myc: 67 kDa.

Positive Controls: C6 whole cell lysate: sc-364373, SH-SY5Y cell lysate: sc-3812 or BJAB whole cell lysate: sc-2207.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



N-Myc (C-4): sc-515099. Western blot analysis of N-Myc expression in Y79 (A), SJRH30 (B) and C6 (C) whole cell lysates.



N-Myc (C-4): sc-515099. Western blot analysis of N-Myc expression in SH-SY5Y (A) and BJAB (B) whole cell lysates.

SELECT PRODUCT CITATIONS

- Hiraiwa, T., et al. 2019. Activation of cell migration via morphological changes in focal adhesions depends on shear stress in MYCN-amplified neuroblastoma cells. J. R. Soc. Interface 16: 20180934.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **N-Myc (B8.4.B): sc-53993** for N-Myc antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647.