

METTL7B (E-10): sc-515267

BACKGROUND

METTL7B (methyltransferase-like protein 7B) is a 244 amino acid protein belonging to the methyltransferase superfamily. METTL7B is believed to have methyltransferase activity, wherein METTL7B catalyzes the transfer of a methyl group from one compound to another. The gene that encodes METTL7B maps to chromosome 12 which makes up about 4.5% of the human genome. A number of skeletal deformities are linked to chromosome 12 including hypochondrogenesis, achondrogenesis and Kniest dysplasia. Chromosome 12 is also home to a homeobox gene cluster which encodes crucial transcription factors for morphogenesis, and the natural killer complex gene cluster encoding C-type lectin proteins which mediate the NK cell response to MHC I interaction. Trisomy 12p leads to facial development defects, seizure disorders and a host of other symptoms varying in severity depending on the extent of mosaicism and is most severe in cases of complete trisomy.

REFERENCES

- Allen, T.L., et al. 1996. Cytogenetic and molecular analysis in trisomy 12p. *Am. J. Med. Genet.* 63: 250-256.
- Yang, W. and Cole, W.G. 1998. Low basal transcripts of the COL2A1 collagen gene from lymphoblasts show alternative splicing of exon 12 in the Kniest form of spondyloepiphyseal dysplasia. *Hum. Mutat.* 1: S1-S2.
- Trowsdale, J., et al. 2001. The genomic context of natural killer receptor extended gene families. *Immunol. Rev.* 181: 20-38.
- Zumkeller, W., et al. 2004. Genotype/phenotype analysis in a patient with pure and complete trisomy 12p. *Am. J. Med. Genet. A* 129A: 261-264.
- Kelley, J., et al. 2005. Comparative genomics of natural killer cell receptor gene clusters. *PLoS Genet.* 1: 129-139.
- Kemmer, L.A., et al. 2006. The natural history of trisomy 12p. *Am. J. Med. Genet. A* 140: 695-703.

CHROMOSOMAL LOCATION

Genetic locus: Mettl7b (mouse) mapping to 10 D3.

SOURCE

METTL7B (E-10) is a mouse monoclonal antibody raised against amino acids 177-244 mapping at the C-terminus of METTL7B of mouse origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

METTL7B (E-10) is available conjugated to agarose (sc-515267 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-515267 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-515267 PE), fluorescein (sc-515267 FITC), Alexa Fluor® 488 (sc-515267 AF488), Alexa Fluor® 546 (sc-515267 AF546), Alexa Fluor® 594 (sc-515267 AF594) or Alexa Fluor® 647 (sc-515267 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-515267 AF680) or Alexa Fluor® 790 (sc-515267 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

METTL7B (E-10) is recommended for detection of METTL7B of mouse and rat origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for METTL7B siRNA (h): sc-96008, METTL7B siRNA (m): sc-149392, METTL7B shRNA Plasmid (h): sc-96008-SH, METTL7B shRNA Plasmid (m): sc-149392-SH, METTL7B shRNA (h) Lentiviral Particles: sc-96008-V and METTL7B shRNA (m) Lentiviral Particles: sc-149392-V.


Molecular Weight of METTL7B: 28 kDa.

Positive Controls: mouse liver extract: sc-2256 or rat liver extract: sc-2395.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



METTL7B (E-10): sc-515267. Western blot analysis of METTL7B expression in mouse liver (A) and rat liver (B) tissue extracts.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.