

G_{β5} (C-6): sc-515379

BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (i.e., a photon, pheromone, odorant, hormone or neurotransmitter), while the effectors (e.g., adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. Each subunit of the G protein complex is encoded by a member of one of three corresponding gene families (α , β , γ). In mammals, there are five different members of the β -subunit family. The β subunits of the G proteins are important regulators of G protein subunits as well as of certain signal transduction receptors and effectors. In contrast to G_{β1-4}, which are at least 83% homologous, G_{β5} is only 50% homologous to the other β subunits. Human G_{β5} is expressed at high levels in brain, pancreas, kidney, and heart.

REFERENCES

- Blatt, C., et al. 1988. Chromosomal localization of genes encoding guanine nucleotide-binding protein subunits in mouse and human. Proc. Natl. Acad. Sci. USA 85: 7642-7646.
- Gautam, N., et al. 1990. G protein diversity is increased by associations with a variety of γ subunits. Proc. Natl. Acad. Sci. USA 87: 7973-7977.

CHROMOSOMAL LOCATION

Genetic locus: GNB5 (human) mapping to 15q21.2; Gnb5 (mouse) mapping to 9 D.

SOURCE

G_{β5} (C-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 27-51 near the N-terminus of G_{β5} of human origin.

PRODUCT

Each vial contains 200 μ g IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

G_{β5} (C-6) is available conjugated to agarose (sc-515379 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-515379 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-515379 PE), fluorescein (sc-515379 FITC), Alexa Fluor® 488 (sc-515379 AF488), Alexa Fluor® 546 (sc-515379 AF546), Alexa Fluor® 594 (sc-515379 AF594) or Alexa Fluor® 647 (sc-515379 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-515379 AF680) or Alexa Fluor® 790 (sc-515379 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-515379 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

G_{β5} (C-6) is recommended for detection of G_{β5} of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for G_{β5} siRNA (h): sc-41770, G_{β5} siRNA (m): sc-41771, G_{β5} shRNA Plasmid (h): sc-41770-SH, G_{β5} shRNA Plasmid (m): sc-41771-SH, G_{β5} shRNA (h) Lentiviral Particles: sc-41770-V and G_{β5} shRNA (m) Lentiviral Particles: sc-41771-V.


Molecular Weight of G_{β5}: 39 kDa.

Positive Controls: mouse brain extract: sc-2253 or human brain extract: sc-364375.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



G_{β5} (C-6): sc-515379. Western blot analysis of G_{β5} expression in mouse brain (A) and human brain (B) tissue extracts.

SELECT PRODUCT CITATIONS

- Zaccone, G., et al. 2020. Expression of Acetylcholine- and G protein-coupled Muscarinic receptor in the Neuroepithelial cells (NECs) of the obligate air-breathing fish, *Arapaima gigas* (Arapaimidae: Teleostei). Zoology 139: 125755.
- Walters, D., et al. 2020. Transcriptome analysis in mice treated with vigabatrin identifies dysregulation of genes associated with retinal signaling circuitry. Epilepsy Res. 166: 106395.

RESEARCH USE

For research use only, not for use in diagnostic procedures.