

GATA-5 (H-6): sc-515422

BACKGROUND

Members of the GATA family share a conserved zinc finger DNA-binding domain and are capable of binding the WGATAR consensus sequence. GATA-1 is erythroid-specific and is responsible for the regulated transcription of erythroid genes. It is an essential component in the generation of the erythroid lineage. GATA-2 is expressed in embryonic brain and liver, HeLa and endothelial cells, as well as in erythroid cells. Studies with a modified GATA consensus sequence, AGATCTTA, have shown that GATA-2 and GATA-3 recognize this mutated consensus while GATA-1 has poor recognition of this sequence. This indicates broader regulatory capabilities of GATA-2 and GATA-3 than GATA-1. GATA-3 is highly expressed in T lymphocytes. GATA-4, GATA-5 and GATA-6 comprise a subfamily of transcription factors. Both GATA-4 and GATA-6 are found in heart, pancreas and ovary; lung and liver tissues exhibit GATA-6, but not GATA-4 expression. GATA-5 expression has been observed in differentiated heart and gut tissues and is present throughout the course of development in the heart. Although expression patterns of the various GATA transcription factors may overlap, it is not yet apparent how the GATA factors are able to discriminate in binding their appropriate target sites.

REFERENCES

1. Ko, L.J., et al. 1991. Murine and human T-lymphocyte GATA-3 factors mediate transcription through a *cis*-regulatory element within the human T cell receptor δ gene enhancer. *Mol. Cell. Biol.* 11: 2778-2784.
2. Dorfman, D.M., et al. 1992. Human transcription factor GATA-2. Evidence for regulation of preproendothelin-1 gene expression in endothelial cells. *J. Biol. Chem.* 267: 1279-1285.
3. Ko, L.J., et al. 1993. DNA-binding specificities of the GATA transcription factor family. *Mol. Cell. Biol.* 13: 4011-4022.
4. Laverriere, A.C., et al. 1994. GATA-4/5/6, a subfamily of three transcription factors transcribed in developing heart and gut. *J. Biol. Chem.* 269: 23177-23184.

CHROMOSOMAL LOCATION

Genetic locus: GATA5 (human) mapping to 20q13.33.

SOURCE

GATA-5 (H-6) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 373-389 at the C-terminus of GATA-5 of human origin.

PRODUCT

Each vial contains 200 μ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-515422 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-515422 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

GATA-5 (H-6) is recommended for detection of GATA-5 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GATA-5 siRNA (h): sc-35456, GATA-5 shRNA Plasmid (h): sc-35456-SH and GATA-5 shRNA (h) Lentiviral Particles: sc-35456-V.

GATA-5 (H-6) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of GATA-5: 45 kDa.

Positive Controls: FHs 173We cell lysate: sc-2417 or HT-1080 whole cell lysate: sc-364183.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



GATA-5 (H-6): sc-515422. Western blot analysis of GATA-5 expression in HT-1080 (A) and FHs 173We (B) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Liu, J., et al. 2022. GATA binding protein 5-mediated transcriptional activation of transmembrane protein 100 suppresses cell proliferation, migration and epithelial-to-mesenchymal transition in prostate cancer DU145 cells. *Bioengineered* 13: 7972-7983.

RESEARCH USE

For research use only, not for use in diagnostic procedures.