

galectin-3 (F-8): sc-515582



The Power to Question

BACKGROUND

Galectins are a family of soluble β -galactoside-binding animal lectins that modulate cell-to-cell adhesion and cell-to-extracellular matrix (ECM) interactions and play a role in tumor progression, pre-mRNA splicing and apoptosis. The galectin-3 protein, also known as Mac-2, hMac-2, GALBP, CBP35 or LGALS3, contains a single carbohydrate binding domain, which binds galactose-containing glycoconjugates. Galectin-3 is expressed in colonic and intestinal epithelium, inflammatory macrophages, papillary and follicular carcinomas, neoplastic astrocytes and some B and T lymphocytes. Upregulated expression of galectin-3 is involved in cancer progression and metastasis. Galectin-3 mediates the endocytosis of β 1 Integrins in a lactose-dependent manner and is associated with thyroid malignancy and Crohn's disease. It may also be used as a marker for diagnosing cases involving Hurthle cell adenomas and carcinomas.

REFERENCES

- Huflejt, M.E., et al. 1997. Strikingly different localization of galectin-3 and galectin-4 in human colon adenocarcinoma T84 cells. Galectin-4 is localized at sites of cell adhesion. *J. Biol. Chem.* 272: 14294-14303.
- Shimonishi, T., et al. 2001. Expression of endogenous galectin-1 and galectin-3 in intrahepatic cholangiocarcinoma. *Hum. Pathol.* 32: 302-310.
- Guittaut, M., et al. 2001. Identification of an internal gene to the human Galectin-3. *J. Biol. Chem.* 276: 2652-2667.
- Jensen-Jarolim, E., et al. 2001. Anti-Galectin-3 IgG autoantibodies in patients with Crohn's disease characterized by means of phage display peptide libraries. *J. Clin. Immunol.* 21: 348-356.

CHROMOSOMAL LOCATION

Genetic locus: LGALS3 (human) mapping to 14q22.3; Lgals3 (mouse) mapping to 14 C1.

SOURCE

galectin-3 (F-8) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-25 at the N-terminus of galectin-3 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-515582 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

galectin-3 (F-8) is recommended for detection of galectin-3 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for galectin-3 siRNA (h): sc-155994, galectin-3 siRNA (m): sc-35443, galectin-3 shRNA Plasmid (h): sc-155994-SH, galectin-3 shRNA Plasmid (m): sc-35443-SH, galectin-3 shRNA (h) Lentiviral Particles: sc-155994-V and galectin-3 shRNA (m) Lentiviral Particles: sc-35443-V.


Molecular Weight of galectin-3: 31 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210, RAW 264.7 whole cell lysate: sc-2211 or C3H/10T1/2 cell lysate: sc-3801.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



galectin-3 (F-8): sc-515582. Western blot analysis of galectin-3 expression in NIH/3T3 (A), RAW 264.7 (B) and C3H/10T1/2 (C) whole cell lysates.



galectin-3 (F-8): sc-515582. Western blot analysis of galectin-3 expression in HeLa (A) and SW480 (B) nuclear extracts.

SELECT PRODUCT CITATIONS

- Schmidt, F., et al. 2020. Flotillin-dependent membrane microdomains are required for functional phagolysosomes against fungal infections. *Cell Rep.* 32: 108017.



See **galectin-3 (B2C10): sc-32790** for galectin-3 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.