

PDLIM5 (G-2): sc-515621

BACKGROUND

PDLIM5 (PDZ and LIM domain 5), also known as L9, ENH (Enigma homolog), LIM or ENH1, is a member of the Enigma family of proteins. Proteins belonging to this family contain an N-terminal PDZ (post-synaptic density-95/discs large/zone occludens-1) domain and one to three C-terminal LIM domains that typically associate with various isoforms of PKC (protein kinase C). Expressed in a wide variety of tissues, PDLIM5 contains three LIM zinc-binding domains and one PDZ domain. In the brain, PDLIM5 colocalizes with synaptic vesicles of neurotransmitters and regulates neuronal calcium signaling through an interaction with PKC ϵ and N-type Ca^{++} CP $\alpha 1\text{B}$ (N-type calcium channel $\alpha 1\text{B}$ subunit). Expression of PDLIM5 is often increased in the brain of patients with schizophrenia, major depression and bipolar disorder. This suggests that PDLIM5 may contribute to the genetic susceptibility of such conditions.

REFERENCES

1. Kuroda, S., et al. 1996. Protein-protein interaction of zinc finger LIM domains with protein kinase C. *J. Biol. Chem.* 271: 31029-31032.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605904. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Wu, M., et al. 2004. Cloning and identification of a novel human gene PDLIM5, a homolog of AD-associated neuronal thread protein (AD7c-NTP). *DNA Seq.* 15: 144-147.
4. Kato, T., et al. 2005. Gene expression and association analyses of LIM (PDLIM5) in bipolar disorder and schizophrenia. *Mol. Psychiatry* 10: 1045-1055.
5. Iga, J., et al. 2006. Gene expression and association analysis of LIM (PDLIM5) in major depression. *Neurosci. Lett.* 400: 203-207.
6. Horiuchi, Y., et al. 2006. A polymorphism in the PDLIM5 gene associated with gene expression and schizophrenia. *Biol. Psychiatry* 59: 434-439.

CHROMOSOMAL LOCATION

Genetic locus: PDLIM5 (human) mapping to 4q22.3.

SOURCE

PDLIM5 (G-2) is a mouse monoclonal antibody raised against amino acids 297-596 mapping at the C-terminus of PDLIM5 of human origin.

PRODUCT

Each vial contains 200 μg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

PDLIM5 (G-2) is available conjugated to agarose (sc-515621 AC), 500 μg /0.25 ml agarose in 1 ml, for IP; to HRP (sc-515621 HRP), 200 μg /ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-515621 PE), fluorescein (sc-515621 FITC), Alexa Fluor® 488 (sc-515621 AF488), Alexa Fluor® 546 (sc-515621 AF546), Alexa Fluor® 594 (sc-515621 AF594) or Alexa Fluor® 647 (sc-515621 AF647), 200 μg /ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-515621 AF680) or Alexa Fluor® 790 (sc-515621 AF790), 200 μg /ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

PDLIM5 (G-2) is recommended for detection of PDLIM5 of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PDLIM5 siRNA (h): sc-76097, PDLIM5 shRNA Plasmid (h): sc-76097-SH and PDLIM5 shRNA (h) Lentiviral Particles: sc-76097-V.

Molecular Weight of PDLIM5: 64 kDa.


Positive Controls: K-562 whole cell lysate: sc-2203 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



PDLIM5 (G-2): sc-515621. Western blot analysis of PDLIM5 expression in K-562 (A) and Hep G2 (B) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Kobayashi, Y., et al. 2021. Ciliary GPCR-based transcriptome as a key regulator of cilia length control. *FASEB Bioadv.* 3: 744-767.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA