

PP2A-B56- β (D-10): sc-515681

BACKGROUND

In eukaryotes, the phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the protein phosphatases. In general, the protein phosphatase (PP) holoenzyme is a trimeric complex composed of a regulatory subunit, a variable subunit, and a catalytic subunit. Four major families of protein phosphatase catalytic subunits have been identified, designated PP1, PP2A, PP2B (calcineurin) and PP2C. An additional protein phosphatase catalytic subunit, PPX (also known as PP4) is a putative member of a novel PP family. The PP2A family comprises subfamily members PP2A α and PP2A β . The PP2A catalytic subunit associates with a variety of regulatory subunits. Regulatory subunits include PP2A-A- α and -A- β , PP2A-B- α and -B- β , PP2A-C- α and -C- β , PP2A-B56- α , -B56- β , -B56- γ and -B56- δ .

REFERENCES

- Ueki, K., et al. 1992. Structure and expression of two isoforms of the murine calmodulin-dependent protein phosphatase regulatory subunit (calcineurin B). *Biochem. Biophys. Res. Commun.* 187: 537-543.
- Cohen, P.T. 1993. Important roles for novel protein phosphatases dephosphorylating serine and threonine residues. *Biochem. Soc. Trans.* 21: 884-888.
- Hendrix, P., et al. 1993. Structure and expression of a 72-kDa regulatory subunit of protein phosphatase 2A. Evidence for different size forms produced by alternative splicing. *J. Biol. Chem.* 268: 15267-15276.
- Mumby, M.C., et al. 1993. Protein serine/threonine phosphatases: structure, regulation, and functions in cell growth. *Physiol. Rev.* 73: 673-699.
- Okubo, S., et al. 1994. A regulatory subunit of smooth muscle myosin bound phosphatase. *Biochem. Biophys. Res. Commun.* 200: 429-434.
- Wera, S., et al. 1995. Serine/threonine protein phosphatases. *Biochem. J.* 311: 17-29.

CHROMOSOMAL LOCATION

Genetic locus: PPP2R5B (human) mapping to 11q13.1; Ppp2r5b (mouse) mapping to 19 A.

SOURCE

PP2A-B56- β (D-10) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 451-472 near the C-terminus of PP2A-B56- β of human origin.

PRODUCT

Each vial contains 200 μ g IgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4°C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

PP2A-B56- β (D-10) is recommended for detection of PP2A-B56- β of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PP2A-B56- β siRNA (h): sc-39183, PP2A-B56- β siRNA (m): sc-39184, PP2A-B56- β shRNA Plasmid (h): sc-39183-SH, PP2A-B56- β shRNA Plasmid (m): sc-39184-SH, PP2A-B56- β shRNA (h) Lentiviral Particles: sc-39183-V and PP2A-B56- β shRNA (m) Lentiviral Particles: sc-39184-V.


Positive Controls: Jurkat whole cell lysate: sc-2204, HeLa whole cell lysate: sc-2200 or IMR-32 cell lysate: sc-2409.

RECOMMENDED SUPPORT REAGENTS


To ensure optimal results, the following support reagents are recommended:

- 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.
- 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml).
- 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.
- 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohisto-mount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



PP2A-B56- β (D-10): sc-515681. Western blot analysis of PP2A-B56- β expression in KNRK (**A**), Jurkat (**B**), HeLa (**C**), IMR-32 (**D**) and OVCAR-3 (**E**) whole cell lysates.



PP2A-B56- β (D-10): sc-515681. Immunoperoxidase staining of formalin fixed, paraffin-embedded human cerebellum tissue showing cytoplasmic staining of Purkinje cells and nuclear staining of a subset of cells in molecular layer (**A**), and of rat brain tissue showing cytoplasmic staining of neuronal cells, glial cells and endothelial cells and neuropil staining (**B**). Blocked with 0.25X UltraCruz® Blocking Reagent: sc-516214. Detection reagents used: m-IgG κ BP-B: sc-516142 and ImmunoCruz® ABC Kit: sc-516216.

RESEARCH USE

For research use only, not for use in diagnostic procedures.