


TRPC6 (B-10): sc-515837



The Power to Question

BACKGROUND

Transient receptor potential cation (TRPC) channels are a superfamily of six transmembrane segment-spanning, gated cation channels. TRPC subtypes mediate store-operated Ca^{2+} entry, a process involving Ca^{2+} influx and replenishment of Ca^{2+} stores formerly emptied through the action of inositol 1,4,5-trisphosphate production and other Ca^{2+} mobilizing agents. TRPC ion channels influence calcium-depletion induced calcium influx processes in response to chemo-, mechano- and osmoregulatory events. Human TRPC6 protein is a 931 amino acid cation channel that is predominantly expressed in placenta, spleen, lung, small intestine and ovary. Activated by diacylglycerol (DAG), TRPC6 comprises the α_1 -adrenoceptor-activated Ca^{2+} -permeable cation channel. The gene encoding human TRPC6 maps to chromosome 11q22.1.

CHROMOSOMAL LOCATION

Genetic locus: TRPC6 (human) mapping to 11q22.1; Trpc6 (mouse) mapping to 9 A1.

SOURCE

TRPC6 (B-10) is a mouse monoclonal antibody raised against amino acids 546-590 mapping within an extracellular domain of TRPC6 of human origin.

PRODUCT

Each vial contains 200 μg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

TRPC6 (B-10) is available conjugated to agarose (sc-515837 AC), 500 μg /0.25 ml agarose in 1 ml, for IP; to HRP (sc-515837 HRP), 200 μg /ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-515837 PE), fluorescein (sc-515837 FITC), Alexa Fluor® 488 (sc-515837 AF488), Alexa Fluor® 546 (sc-515837 AF546), Alexa Fluor® 594 (sc-515837 AF594) or Alexa Fluor® 647 (sc-515837 AF647), 200 μg /ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-515837 AF680) or Alexa Fluor® 790 (sc-515837 AF790), 200 μg /ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

TRPC6 (B-10) is recommended for detection of TRPC6 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for TRPC6 siRNA (h): sc-42672, TRPC6 siRNA (m): sc-42673, TRPC6 shRNA Plasmid (h): sc-42672-SH, TRPC6 shRNA Plasmid (m): sc-42673-SH, TRPC6 shRNA (h) Lentiviral Particles: sc-42672-V and TRPC6 shRNA (m) Lentiviral Particles: sc-42673-V.

Positive Controls: A549 cell lysate: sc-2413, WI-38 whole cell lysate: sc-364260 or BT-20 cell lysate: sc-2223.


STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



TRPC6 (B-10): sc-515837. Western blot analysis of TRPC6 expression in A549 (A), DU 145 (B), F9 (C) and AT3B-1 (D) whole cell lysates.



TRPC6 (B-10): sc-515837. Western blot analysis of TRPC6 expression in WI-38 (A), BT-20 (B) and RT-4 (C) whole cell lysates.

SELECT PRODUCT CITATIONS

- Dong, F., et al. 2019. Chrysin alleviates chronic hypoxia-induced pulmonary hypertension by reducing intracellular calcium concentration in pulmonary arterial smooth muscle cells. *J. Cardiovasc. Pharmacol.* 74: 426-435.
- Imnaten, M., et al. 2020. Receptor potential channels TRPC1/TRPC6 regulate lamina cribrosa cell extracellular matrix gene transcription and proliferation. *Exp. Eye Res.* 193: 107980.
- Castillo-Galán, S., et al. 2020. Stim-activated TRPC-ORAI channels in pulmonary hypertension induced by chronic intermittent hypoxia. *Pulm. Circ.* 10: 13-22.
- Chen, L., et al. 2021. Atractylodis rhizoma water extract attenuates fructose-induced glomerular injury in rats through anti-oxidation to inhibit TRPC6/p-CaMK4 signaling. *Phytomedicine* 91: 153643.
- Chen, L., et al. 2021. Atractylodin inhibits fructose-induced human podocyte hypermotility via anti-oxidant to down-regulate TRPC6/p-CaMK4 signaling. *Eur. J. Pharmacol.* 913: 174616.
- Wong, C.J.K., et al. 2022. Brief exposure to directionally-specific pulsed electromagnetic fields stimulates extracellular vesicle release and is antagonized by streptomycin: a potential regenerative medicine and food industry paradigm. *Biomaterials* 287: 121658.
- Lu, T., et al. 2022. TRPC6 N338S is a gain-of-function mutant identified in patient with doxorubicin-induced cardiotoxicity. *Biochim. Biophys. Acta Mol. Basis Dis.* 1868: 166505.
- Cai, J., et al. 2022. Upregulation of TRPC6 inhibits astrocyte activation and proliferation after spinal cord injury in rats by suppressing AQP4 expression. *Brain Res. Bull.* 190: 12-21.
- Wang, M., et al. 2023. TRPC6 deletion enhances eNOS expression and reduces LPS-induced acute lung injury. *Int. J. Mol. Sci.* 24: 16756.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.